

Twisted Hamiltonian Lie Algebras and Their Multiplicity-Free Representations ^{*}

Ling Chen [†]

Institute of Applied Mathematics, Academy of Mathematics & System Sciences,
Chinese Academy of Sciences, Beijing 100190, China

Abstract

We construct a class of new Lie algebras by generalizing the one-variable Lie algebras generated by the quadratic conformal algebras (or corresponding Hamiltonian operators) associated to Poisson algebras and a quasi-derivation found by Xu. These algebras can be viewed as certain twists of Xu's generalized Hamiltonian Lie algebras. The simplicity of these algebras is completely determined. Moreover, we construct a family of multiplicity-free representations of these Lie algebras and prove their irreducibility.

Key words: Hamiltonian Lie algebras, representation, simplicity, irreducibility.

1 Introduction

Simple Lie algebras of Cartan type play important roles in the structure theory of Lie algebras. They also have many applications in the areas of geometry and mathematical physics. After Kac [2] gave an abstract definition of Lie algebras of Cartan type by derivations, various graded generalizations were introduced and studied by Kawamoto [3], Osborn [6], Dokovic and Zhao [1], and Zhao [25]. Motivated from his works on quadratic conformal algebras [23] and Hamiltonian operators [24], Xu [22] found certain nongraded generalizations of the Lie algebras of Cartan type and determined their simplicity. Indeed, Xu's algebras in general do not contain any toral Cartan subalgebras, so traditional approaches to simple Lie algebras do not work in his case. In particular, Xu's generalizations

^{*}2000 Mathematics Subject Classification. Primary 17B10, 17B20, 17B65.

[†]Corresponding author. E-mail: chenling@amss.ac.cn (L. Chen)

of Hamiltonian Lie algebras were constructed from a skew-symmetric $\mathbb{Z}$ -bilinear form and an even number of locally-finite commuting derivations. Hamiltonian Lie algebras play key roles in both classical and quantum physics. They are also fundamental algebraic structure in symplectic geometry.

Passman [8] gave a certain necessary and sufficient condition on derivations for a Lie algebra of generalized Witt type to be simple. The isomorphic structural spaces of Xu's nongraded Lie algebras of Cartan type were determined in [15–18]. Moreover, Xu [21] also introduced certain nongraded generalizations of the Lie algebras of Block type and determined their simplicity.

On the representation side, Shen [11–13] introduced mixed product of graded modules over graded Lie algebras of Cartan type (also known as Larsson functor (cf. [4])) and obtained certain irreducible modules. Rao [9, 10] constructed some irreducible weight modules over the derivation Lie algebra of the algebra of Laurent polynomials based on Shen's mixed product. Lin and Tan [5] did the similar thing over the derivation Lie algebra of the algebra of quantum torus. Zhao [27] determined the module structure of Shen's mixed product over Xu's nongraded Lie algebras of Witt type.

Multiplicity-free representations are important objects in Lie theory due to their neat formulae and natural applications. Su and Zhao [20] classified the multiplicity-free representations over generalized Virasoro algebras. Moreover, Zhao [26] gave a classification of the multiplicity-free representations over graded generalized Witt algebras. Furthermore, Su and Zhou [19] classified the analogues of multiplicity-free representations over Xu's nongraded generalized Witt algebras.

In [23], Xu classified certain quadratic conformal algebras and constructed such algebras from Poisson algebras and a quasi-derivation. Conformal algebras are the local structures of certain one-variable Lie algebras. They are also equivalent to certain linear Hamiltonian operators (cf. [24]). We construct a class of new Lie algebras by generalizing the one-variable Lie algebras generated by the quadratic conformal algebras associated to Poisson algebras and a quasi-derivation. These algebras can be viewed as certain twists of Xu's generalized Hamiltonian Lie algebras. The simplicity of these algebras is completely determined. Moreover, we construct a family of multiplicity-free representations of these Lie algebras and prove their irreducibility. Below we give more technical details.

Throughout this paper, let $\mathbb{F}$ be a field with characteristic 0. All the vector spaces are assumed over $\mathbb{F}$. Let $(\mathcal{A}, [\cdot, \cdot], \cdot)$ be a Poisson algebra, that is, $(\mathcal{A}, [\cdot, \cdot])$ forms a Lie algebra, $(\mathcal{A}, \cdot)$ forms a commutative associative algebra and the following compatibility condition holds:

$$[u, v \cdot w] = [u, v] \cdot w + v \cdot [u, w] \quad \text{for } u, v, w \in \mathcal{A}. \quad (1.1)$$

Let ∂_1 and ∂_2 be two commuting derivations of $(\mathcal{A}, \cdot)$. Suppose that ∂_2 is also a derivation of the Lie algebra $(\mathcal{A}, [\cdot, \cdot])$ but ∂_1 is a quasi-derivation of $(\mathcal{A}, [\cdot, \cdot])$ in the following sense:

$$\partial_1([u, v]) = [\partial_1(u), v] + [u, \partial_1(v)] + c[u, v] \quad \text{for } u, v \in \mathcal{A}, \quad (1.2)$$

where $c \in \mathbb{F}$ is a fixed constant. Define a new algebraic operation $[\cdot, \cdot]_1$ on $\mathcal{A}$ by

$$[u, v]_1 = [u, v] + (\partial_1 + c)(u) \cdot \partial_2(v) - \partial_2(u) \cdot (\partial_1 + c)(v) \quad \text{for } u, v \in \mathcal{A}. \quad (1.3)$$

It can be verified that $(\mathcal{A}, [\cdot, \cdot]_1)$ also forms a Lie algebra, which is motivated from the one-variable Lie algebras generated by the quadratic conformal algebras associated to Poisson algebras and a quasi-derivation in [23]. If $[\cdot, \cdot]$ is the trivial Lie bracket, that is, $[u, v] = 0$ for $u, v \in \mathcal{A}$, then $(\mathcal{A}, [\cdot, \cdot]_1)$ with $c = 1$ is exactly Xu's generalization of Block algebra in [21]. In the case $c = 0$, (1.3) can be viewed as the iterated construction of Hamiltonian Lie algebras. When $\mathcal{A} = \mathbb{F}[x_1, \dots, x_{2n+1}]$, $\partial_2 = \partial_{x_{2n+1}}$, ∂_1 is the grading operator on $\mathcal{A}$ with respect to $\{x_1, \dots, x_{2n}\}$ and

$$[f, g] = \sum_{i=1}^n (\partial_{x_i}(f) \partial_{x_{n+i}}(g) - \partial_{x_{n+i}}(f) \partial_{x_i}(g)) \quad \text{for } f, g \in \mathcal{A}, \quad (1.4)$$

the Lie algebra $(\mathcal{A}, [\cdot, \cdot]_1)$ with $c = -2$ is exactly the canonical Lie algebra of Contact type. In this paper, we will deal with another case, which can be viewed as a generalization of Xu's four-derivation analogues of Block algebras in [21].

Denote by $\mathbb{Z}$ the ring of integers and by $\mathbb{N}$ the additive semigroup of nonnegative integers. For two integers $i \leq j$, we denote $\overline{i, j} = \{i, i+1, \dots, j\}$. Let n be a positive integer. Picking

$$\mathcal{J}_p \in \{\{0\}, \mathbb{N}\} \quad \text{for } p \in \overline{1, 2n}, \quad (1.5)$$

we have the following product of additive semigroups

$$\mathcal{J} = \mathcal{J}_1 \times \mathcal{J}_2 \times \dots \times \mathcal{J}_{2n}. \quad (1.6)$$

Let Γ be a nonzero torsion-free abelian group, which is also viewed as a $\mathbb{Z}$ -module throughout this paper. Let $\phi(\cdot, \cdot) : \Gamma \times \Gamma \rightarrow \mathbb{F}$ be a skew-symmetric $\mathbb{Z}$ -bilinear form and let $\{\varphi_p : \Gamma \rightarrow \mathbb{F} \mid p \in \overline{1, 2n}\}$ be $2n$ additive group homomorphisms such that

$$\varphi_p(\Gamma) + \mathcal{J}_p \neq \{0\} \quad \text{for } p \in \overline{1, 2n}, \quad (1.7)$$

$$\text{Rad}_\phi \cap \left(\bigcap_{p \in \overline{1, 2n}} \ker \varphi_p \right) = \{0\}, \quad (1.8)$$

and

$$\text{Rad}_\phi \cap \left(\bigcap_{q \neq p \in \overline{1, 2n}} \ker_{\varphi_p} \right) \setminus \ker_{\varphi_q} \neq \emptyset, \quad \text{if } \varphi_q \neq 0, \quad q \in \overline{1, 2n}. \quad (1.9)$$

We choose fixed elements

$$\sigma_{2s+1} = \sigma_{2s+2} \in \text{Rad}_\phi \cap \left(\bigcap_{2s+1, 2s+2 \neq p \in \overline{1, 2n}} \ker_{\varphi_p} \right) \quad (1.10)$$

such that $\sigma_{2s+1} \neq 0$ if $\varphi_{2s+1} \neq 0$ or $\varphi_{2s+2} \neq 0$ for $s \in \overline{1, n-1}$.

Let $\mathcal{A}$ be a vector space with a basis $\{x^{\alpha, \mathbf{i}} \mid (\alpha, \mathbf{i}) \in \Gamma \times \mathcal{J}\}$. Define an algebraic operation $\cdot$ on $\mathcal{A}$ by

$$x^{\alpha, \mathbf{i}} \cdot x^{\beta, \mathbf{j}} = x^{\alpha+\beta, \mathbf{i}+\mathbf{j}} \quad \text{for } (\alpha, \mathbf{i}), (\beta, \mathbf{j}) \in \Gamma \times \mathcal{J}. \quad (1.11)$$

Then $\mathcal{A}$ forms a commutative associative algebra with an identity element $x^{0, \mathbf{0}}$, which will simply be denoted by 1 in the rest of this paper. Throughout this paper, the notion “.” will be invisible in a product when the context is clear.

Use $\mathbf{i} = (i_1, i_2, \dots, i_{2n})$ to denote an element in $\mathcal{J}$. Moreover, we denote

$$a_{[p]} = (0, \dots, \overset{p}{a}, 0, \dots, 0), \quad a \in \mathcal{J}_p. \quad (1.12)$$

For $p \in \overline{1, 2n}$, we define $\partial_p \in \text{End } \mathcal{A}$ by

$$\partial_p(x^{\alpha, \mathbf{i}}) = \varphi_p(\alpha)x^{\alpha, \mathbf{i}} + i_p x^{\alpha, \mathbf{i}-1_{[p]}} \quad (1.13)$$

for $(\alpha, \mathbf{i}) \in \Gamma \times \mathcal{J}$, where we adopt the convention that if a notion is not defined but technically appears in an expression, we always treat it as zero; for instance, $x^{\alpha, -1_{[1]}} = 0$ for any $\alpha \in \Gamma$. It can be verified that $\{\partial_p \mid p \in \overline{1, 2n}\}$ are commuting derivations of $\mathcal{A}$. For any $\alpha \in \Gamma$, we set $\mathcal{A}_\alpha = \text{Span}\{x^{\alpha, \mathbf{j}} \mid \mathbf{j} \in \mathcal{J}\}$. Assume that $\text{Rad}_\phi \cap (\bigcap_{p=2}^{2n} \ker_{\varphi_p})$ has an element ε such that $\varphi_1(\varepsilon) = 1$. Then we have a Lie bracket $[\cdot, \cdot]$ of the type (1.3) on $\mathcal{A}$ determined by

$$\begin{aligned} [u, w] &= x^{\varepsilon, \mathbf{0}} \left(\sum_{s=1}^{n-1} x^{\sigma_{2s+1}, \mathbf{0}} (\partial_{2s+1}(u) \partial_{2s+2}(w) - \partial_{2s+1}(w) \partial_{2s+2}(u)) + \phi(\alpha, \beta) u w \right) \\ &\quad + (\partial_1 + 1)(u) \partial_2(w) - (\partial_1 + 1)(w) \partial_2(u) \end{aligned} \quad (1.14)$$

for $u \in \mathcal{A}_\alpha$ and $w \in \mathcal{A}_\beta$.

Notice that $x^{-\varepsilon, \mathbf{0}}$ is a central element of $\mathcal{A}$. Form a quotient Lie algebra $\mathcal{H} = \mathcal{A} / \mathbb{F} x^{-\varepsilon, \mathbf{0}}$, whose induced Lie bracket is still denoted by $[\cdot, \cdot]$ when the context is clear. We call the Lie algebra $(\mathcal{H}, [\cdot, \cdot])$ a *twisted Hamiltonian Lie algebra*. Denote $\sigma = \sum_{s=1}^{n-1} \sigma_{2s+1} - 2\varepsilon$.

Theorem 1 *The Lie algebra $(\mathcal{H}, [\cdot, \cdot])$ is simple if $\mathcal{J} \neq \{\mathbf{0}\}$. If $\mathcal{J} = \{\mathbf{0}\}$, then $\mathcal{H}^{(1)} = [\mathcal{H}, \mathcal{H}]$ is a simple Lie algebra and $\mathcal{H} = \mathcal{H}^{(1)} \oplus (\mathbb{F}x^{\sigma, \mathbf{0}} + \mathbb{F}x^{-\varepsilon, \mathbf{0}})$.*

Take $\vec{\xi} = (\xi_1, \dots, \xi_{2n}) \in \mathbb{F}^{2n}$ and an additive group homomorphism $f : \Gamma \rightarrow \mathbb{F}$ such that $\text{Rad}_\phi \subset \ker f$. For $u \in \mathcal{A}$, we denote $\bar{u} = u + \mathbb{F}x^{-\varepsilon, \mathbf{0}}$. In particular, we denote $\bar{x}^{\alpha, \mathbf{i}} = x^{\alpha, \mathbf{i}} + \mathbb{F}x^{-\varepsilon, \mathbf{0}}$ for $(\alpha, \mathbf{i}) \in \Gamma \times \mathcal{J}$. It can be verified that $\mathcal{A}$ forms a *multiplicity-free* $\mathcal{H}$ -module with the action determined by

$$\begin{aligned} \bar{u}.v &= x^{\varepsilon, \mathbf{0}} \left(\sum_{s=1}^{n-1} x^{\sigma_{2s+1}, \mathbf{0}} (\partial_{2s+1}(u)(\partial_{2s+2} + \xi_{2s+2})(v) - \partial_{2s+2}(u)(\partial_{2s+1} + \xi_{2s+1})(v)) \right. \\ &\quad \left. + (\phi(\alpha, \beta) + f(\alpha))uv \right) + (\partial_1 + 1)(u)(\partial_2 + \xi_2)(v) - \partial_2(u)(\partial_1 + \xi_1)(v) \end{aligned} \quad (1.15)$$

for $u \in \mathcal{A}_\alpha$ and $v \in \mathcal{A}_\beta$. To make it distinguishable, we denote by $\mathcal{A}_{\vec{\xi}, f}$ the module $\mathcal{A}$ with the above action. In particular, denoting $\vec{\xi}_0 = (0, \dots, 0, 1) \in \mathbb{F}^{2n}$, module $\mathcal{A}_{\vec{\xi}_0, 0}$ is induced from the adjoint representation of the Lie algebra $\mathcal{A}$. Moreover, it is straightforward to verify that $\mathcal{A}_{\vec{\xi}', f'} \cong \mathcal{A}_{\vec{\xi}, f}$ if there exists $\mu' \in \Gamma$ such that $\xi'_p - \xi_p = \varphi_p(\mu')$ for $p \in \overline{1, 2n}$, and $f'(\beta) - f(\beta) = \phi(\beta, \mu')$ for any $\beta \in \Gamma$.

Theorem 2 *If there exists $\mu \in \Gamma$ such that*

$$\xi_p = \varphi_p(\mu) \quad \text{for } p \in \overline{1, 2n}, \quad (1.16)$$

and

$$f(\alpha) = \phi(\alpha, \mu) \quad \text{for any } \alpha \in \Gamma, \quad (1.17)$$

then $\mathcal{A}_{\vec{\xi}, f} \cong \mathcal{A}_{\vec{\xi}_0, 0}$. Otherwise, $\mathcal{A}_{\vec{\xi}, f}$ is irreducible.

We shall present the proofs of Theorem 1 and Theorem 2 in sections 2 and 3, respectively.

2 Proof of Theorem 1

In this section, we will determine the simplicity of the twisted Hamiltonian Lie algebra $\mathcal{H}$. In order to prove simplicity, we need the following Lemma.

Lemma 3 *Let $\mathbf{T}$ be a linear transformation on a vector space U , and let U_1 be a subspace of U such that $\mathbf{T}(U_1) \subset U_1$. Suppose that $u_1, u_2, \dots, u_n$ are eigenvectors of $\mathbf{T}$ corresponding to different eigenvalues. If $\sum_{j=1}^n u_j \in U_1$, then $u_1, u_2, \dots, u_n \in U_1$.*

Let $\mathcal{I}$ be any ideal of $\mathcal{A}$ that strictly contains $\mathbb{F}x^{-\varepsilon, \mathbf{0}}$. When $\mathcal{J} \neq \{\mathbf{0}\}$, proving the simplicity of $(\mathcal{H}, [\cdot, \cdot])$ is equivalent to proving that $\mathcal{I} = \mathcal{A}$.

Our proof of Theorem 1 is organized as follows. Firstly, we will determine some preliminary facts about the ideal $\mathcal{I}$. Then, we will process the proof in four steps.

First by (1.14), we have

$$\begin{aligned}
& [x^{\alpha, \mathbf{i}}, x^{\beta, \mathbf{j}}] \\
&= ((\varphi_1(\alpha) + 1)\varphi_2(\beta) - \varphi_2(\alpha)(\varphi_1(\beta) + 1))x^{\alpha+\beta, \mathbf{i}+\mathbf{j}} + (i_1\varphi_2(\beta) - j_1\varphi_2(\alpha)) \cdot \\
&\quad x^{\alpha+\beta, \mathbf{i}+\mathbf{j}-1_{[1]}} + (j_2(\varphi_1(\alpha) + 1) - i_2(\varphi_1(\beta) + 1))x^{\alpha+\beta, \mathbf{i}+\mathbf{j}-1_{[2]}} + (i_1j_2 - i_2j_1) \cdot \\
&\quad x^{\alpha+\beta, \mathbf{i}+\mathbf{j}-1_{[1]}-1_{[2]}} + x^{\varepsilon, \mathbf{0}} \left(\sum_{s=1}^{n-1} x^{\sigma_{2s+1}, \mathbf{0}} ((\varphi_{2s+1}(\alpha)\varphi_{2s+2}(\beta) - \varphi_{2s+2}(\alpha)\varphi_{2s+1}(\beta)) \right. \\
&\quad \cdot x^{\alpha+\beta, \mathbf{i}+\mathbf{j}} + (i_{2s+1}\varphi_{2s+2}(\beta) - j_{2s+1}\varphi_{2s+2}(\alpha))x^{\alpha+\beta, \mathbf{i}+\mathbf{j}-1_{[2s+1]}} + (j_{2s+2}\varphi_{2s+1}(\alpha) \\
&\quad - i_{2s+2}\varphi_{2s+1}(\beta))x^{\alpha+\beta, \mathbf{i}+\mathbf{j}-1_{[2s+2]}} + (i_{2s+1}j_{2s+2} - i_{2s+2}j_{2s+1}) \cdot \\
&\quad \left. x^{\alpha+\beta, \mathbf{i}+\mathbf{j}-1_{[2s+1]}-1_{[2s+2]}} \right) + \phi(\alpha, \beta)x^{\alpha+\beta, \mathbf{i}+\mathbf{j}} \quad (2.1)
\end{aligned}$$

for $(\alpha, \mathbf{i}), (\beta, \mathbf{j}) \in \Gamma \times \mathcal{J}$. In particular,

$$[1, x^{\beta, \mathbf{j}}] = \varphi_2(\beta)x^{\beta, \mathbf{j}} + j_2x^{\beta, \mathbf{j}-1_{[2]}} \quad \text{for } (\beta, \mathbf{j}) \in \Gamma \times \mathcal{J}. \quad (2.2)$$

Set

$$\Gamma' = \{\varphi_2(\alpha) \mid \alpha \in \Gamma\}. \quad (2.3)$$

For $\lambda \in \Gamma'$, we define

$$\mathcal{A}_{(\lambda)} = \{u \in \mathcal{A} \mid (\text{ad}_1 - \lambda)^m(u) = 0 \text{ for some } m \in \mathbb{N}\}. \quad (2.4)$$

Then by (2.2), we have

$$\mathcal{A} = \bigoplus_{\lambda \in \Gamma'} \mathcal{A}_{(\lambda)}. \quad (2.5)$$

In particular,

$$\mathcal{I} = \bigoplus_{\lambda \in \Gamma'} \mathcal{I}_{\lambda}, \quad \mathcal{I}_{\lambda} = \mathcal{I} \cap \mathcal{A}_{(\lambda)}. \quad (2.6)$$

For any $(\beta, \mathbf{j}) \in \Gamma \times \mathcal{J}$, we find

$$\begin{aligned}
[x^{-\beta, \mathbf{0}}, x^{\beta, \mathbf{j}}] &= \sum_{s=1}^{n-1} x^{\varepsilon+\sigma_{2s+1}, \mathbf{0}} (j_{2s+1}\varphi_{2s+2}(\beta)x^{0, \mathbf{j}-1_{[2s+1]}} - j_{2s+2}\varphi_{2s+1}(\beta)x^{0, \mathbf{j}-1_{[2s+2]}}) \\
&\quad + 2\varphi_2(\beta)x^{0, \mathbf{j}} + j_1\varphi_2(\beta)x^{0, \mathbf{j}-1_{[1]}} - j_2(\varphi_1(\beta) - 1)x^{0, \mathbf{j}-1_{[2]}} \quad (2.7)
\end{aligned}$$

by (2.1). If $\mathcal{I}_\lambda \neq \{0\}$ for some $0 \neq \lambda \in \Gamma'$, we pick any

$$0 \neq u = \sum_{(\alpha, \mathbf{i}) \in \Gamma \times \mathcal{J}} d_{\alpha, \mathbf{i}} x^{\alpha, \mathbf{i}} \in \mathcal{I}_\lambda, \quad d_{\alpha, \mathbf{i}} \in \mathbb{F}. \quad (2.8)$$

Assume $d_{\beta, \mathbf{j}} \neq 0$ for some $(\beta, \mathbf{j}) \in \Gamma \times \mathcal{J}$. Denote

$$|\mathbf{i}| = \sum_{p=1}^{2n} i_p \quad \text{for } \mathbf{i} \in \mathcal{J}. \quad (2.9)$$

Fixing β and considering the nonzero terms $d_{\beta, \mathbf{i}} x^{\beta, \mathbf{i}}$ in u with the largest value $|\mathbf{i}|$, we have

$$[x^{-\beta, \mathbf{0}}, u] \in \mathcal{I}_0 \setminus \mathbb{F} x^{-\varepsilon, \mathbf{0}} \quad (2.10)$$

by (2.7) and the fact $\varepsilon \neq \mathbf{0}$. Thus (2.6) implies that $\mathcal{I}_0$ strictly contains $\mathbb{F} x^{-\varepsilon, \mathbf{0}}$.

Observe

$$[1, x^{\beta, \mathbf{j}}] = j_2 x^{\beta, \mathbf{j}-1[2]} \quad \text{for } (\beta, \mathbf{j}) \in \ker_{\varphi_2} \times \mathcal{J} \quad (2.11)$$

and

$$[x^{\varepsilon, \mathbf{0}}, x^{-\varepsilon, 1[2]}] = 2x^{0, \mathbf{0}} \quad (2.12)$$

by (2.1). Set

$$\mathcal{I}_{[0]} = \mathcal{I} \cap \left(\sum_{(\alpha, \mathbf{i}) \in \ker_{\varphi_2} \times \mathcal{J}; i_2=0} \mathbb{F} x^{\alpha, \mathbf{i}} \right). \quad (2.13)$$

By (2.11) and (2.12) and repeatedly acting of ad_1 on $\mathcal{I}_0$, we can prove that

$$\mathcal{I}_{[0]} \text{ strictly contains } \mathbb{F} x^{-\varepsilon, \mathbf{0}}. \quad (2.14)$$

For $k \in \mathbb{N}$, let

$$\mathcal{A}_{\langle k \rangle} = \text{span}\{x^{\alpha, \mathbf{i}} \mid (\alpha, \mathbf{i}) \in \Gamma \times \mathcal{J}, |\mathbf{i}| \leq k\}. \quad (2.15)$$

Then we have

$$\mathcal{A} = \bigcup_{k=0}^{\infty} \mathcal{A}_{\langle k \rangle}. \quad (2.16)$$

Set

$$\hat{k} = \min\{k \in \mathbb{N} \mid (\mathcal{I}_{[0]} \cap \mathcal{A}_{\langle k \rangle}) \setminus \mathbb{F} x^{-\varepsilon, \mathbf{0}} \neq \emptyset\}. \quad (2.17)$$

For any $u \in (\mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle}) \setminus \mathbb{F} x^{-\varepsilon, \mathbf{0}}$, we write it as

$$u = \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\alpha, \mathbf{i}) \in \ker_{\varphi_2} \times \mathcal{J} \\ i_2=0, |\mathbf{i}|=\hat{k}}} a_{\alpha, \mathbf{i}} x^{\alpha, \mathbf{i}} + u', \quad (2.18)$$

where $a_{\alpha, \mathbf{i}} \in \mathbb{F}$, $u' \in \mathcal{A}_{\langle \hat{k}-1 \rangle} + \mathbb{F}x^{-\varepsilon, \mathbf{0}}$. Moreover, we define

$$\iota(u) = |\{\alpha \in \ker_{\varphi_2} \mid a_{\alpha, \mathbf{i}} \neq 0 \text{ for some } \mathbf{i} \in \mathcal{J}, |\mathbf{i}| = \hat{k}\}|. \quad (2.19)$$

By (2.17), $\iota(u) > 0$. Furthermore, we set

$$\iota = \min\{\iota(w) \mid w \in (\mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle}) \setminus \mathbb{F}x^{-\varepsilon, \mathbf{0}}\}. \quad (2.20)$$

Choose $u \in (\mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle}) \setminus \mathbb{F}x^{-\varepsilon, \mathbf{0}}$ such that $\iota(u) = \iota$. Write u as in (2.18). We will process our proof in four steps.

Step 1. $\alpha = \beta$ *whenever* $a_{\alpha, \mathbf{i}} a_{\beta, \mathbf{j}} \neq 0$.

Firstly, we want to prove that

$$\varphi_1(\alpha) = \varphi_1(\beta) \text{ whenever } a_{\alpha, \mathbf{i}} a_{\beta, \mathbf{j}} \neq 0. \quad (2.21)$$

If $\mathcal{J}_2 = \mathbb{N}$, we get

$$\begin{aligned} & \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle} \ni [x^{0, 1[2]}, u] \\ \equiv & \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\alpha, \mathbf{i}) \in \ker_{\varphi_2} \times \mathcal{J} \\ i_2=0, |\mathbf{i}|=\hat{k}}} -a_{\alpha, \mathbf{i}} (\varphi_1(\alpha) + 1) x^{\alpha, \mathbf{i}} \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}}. \end{aligned} \quad (2.22)$$

By the minimality of $\iota(u)$ and Lemma 3, we have

$$\varphi_1(\alpha) + 1 = \varphi_1(\beta) + 1 \text{ whenever } a_{\alpha, \mathbf{i}} a_{\beta, \mathbf{j}} \neq 0, \quad (2.23)$$

that is,

$$\varphi_1(\alpha) = \varphi_1(\beta) \text{ whenever } a_{\alpha, \mathbf{i}} a_{\beta, \mathbf{j}} \neq 0. \quad (2.24)$$

Then (2.21) holds.

If $\mathcal{J}_2 = \{0\}$, then $\varphi_2 \not\equiv 0$ by (1.7). Picking $\tau \in \text{Rad}_{\phi} \cap (\cap_{2 \neq p \in \overline{1, 2n}} \ker_{\varphi_p}) \setminus \ker_{\varphi_2}$, we have

$$\begin{aligned} \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle} & \ni [x^{-\tau, \mathbf{0}}, [x^{\tau, \mathbf{0}}, u]] \equiv \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\alpha, \mathbf{i}) \in \ker_{\varphi_2} \times \mathcal{J} \\ i_2=0, |\mathbf{i}|=\hat{k}}} -a_{\alpha, \mathbf{i}} \varphi_2^2(\tau) (\varphi_1(\alpha) + 1) \cdot \\ & (\varphi_1(\alpha) + 2) x^{\alpha, \mathbf{i}} \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}}. \end{aligned} \quad (2.25)$$

By the minimality of $\iota(u)$ and Lemma 3, we get

$$(\varphi_1(\alpha) + 1)(\varphi_1(\alpha) + 2) = (\varphi_1(\beta) + 1)(\varphi_1(\beta) + 2) \quad (2.26)$$

whenever $a_{\alpha, \mathbf{i}} a_{\beta, \mathbf{j}} \neq 0$, which implies

$$\varphi_1(\alpha) = \varphi_1(\beta) \text{ or } \varphi_1(\alpha + \beta) + 3 = 0 \quad \text{whenever } a_{\alpha, \mathbf{i}} a_{\beta, \mathbf{j}} \neq 0. \quad (2.27)$$

Assume that there exist $\alpha, \beta \in \Gamma$ such that $\varphi_1(\alpha) \neq \varphi_1(\beta)$, $\varphi_1(\alpha + \beta) + 3 = 0$ and $a_{\alpha, \mathbf{i}} a_{\beta, \mathbf{j}} \neq 0$. Without loss of generality, we assume that $\varphi_1(\alpha) + 1 \neq 0$. Note that $\varphi_1 \not\equiv 0$ and $\varphi_2 \not\equiv 0$. According to (1.9), we pick $\tau \in \text{Rad}_\phi \cap (\bigcap_{2 \neq p \in \overline{1, 2n}} \ker_{\varphi_p}) \setminus \ker_{\varphi_2}$ and $\tau' \in \text{Rad}_\phi \cap (\bigcap_{p \in \overline{2, 2n}} \ker_{\varphi_p}) \setminus \ker_{\varphi_1}$ such that

$$\varphi_1(\alpha + \tau') + 1 \neq 0 \text{ and } \varphi_1(\alpha + \tau') + 2 \neq 0. \quad (2.28)$$

Then we find

$$\begin{aligned} \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle} \ni [x^{-\tau, \mathbf{0}}, [x^{\tau + \tau', \mathbf{0}}, u]] &\equiv \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\gamma, \mathbf{l}) \in \ker_{\varphi_2} \times \mathcal{J} \\ l_2 = 0, |\mathbf{l}| = \hat{k}}} -a_{\gamma, \mathbf{l}} \varphi_2^2(\tau) (\varphi_1(\gamma) + 1) \cdot \\ &(\varphi_1(\gamma + \tau') + 2) x^{\gamma + \tau', \mathbf{l}} \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}}. \end{aligned} \quad (2.29)$$

Since $\iota(u)$ is minimal, $\alpha + \tau' \neq -\varepsilon$ due to $\varphi_1(\alpha + \tau') + 1 \neq 0$, and

$$a_{\alpha, \mathbf{i}} \varphi_2^2(\tau) (\varphi_1(\alpha) + 1) (\varphi_1(\alpha + \tau') + 2) \neq 0, \quad (2.30)$$

we have

$$a_{\beta, \mathbf{j}} \varphi_2^2(\tau) (\varphi_1(\beta) + 1) (\varphi_1(\beta + \tau') + 2) \neq 0. \quad (2.31)$$

But

$$\varphi_1(\alpha + \tau') \neq \varphi_1(\beta + \tau'), \quad (2.32)$$

and

$$\varphi_1(\alpha + \tau' + \beta + \tau') + 3 = 2\varphi_1(\tau') \neq 0, \quad (2.33)$$

which contradicts (2.27) if we replace u by $[x^{-\tau, \mathbf{0}}, [x^{\tau + \tau', \mathbf{0}}, u]]$. Thus the first equation in (2.27) holds, and so does (2.21).

Secondly, we want to prove that

$$\varphi_q(\alpha) = \varphi_q(\beta) \quad \text{whenever } a_{\alpha, \mathbf{i}} a_{\beta, \mathbf{j}} \neq 0 \text{ for } q \in \overline{3, 2n}. \quad (2.34)$$

Fixing q , if $\varphi_q \equiv 0$, it is done. For $s \in \overline{0, n-1}$, we define

$$(2s+1)' = 2s+2, \quad (2s+2)' = 2s+1, \quad \epsilon_{2s+1} = 1, \quad \epsilon_{2s+2} = -1. \quad (2.35)$$

Assume $\varphi_q \not\equiv 0$ and $\varphi_{q'} \not\equiv 0$. Then for any $0 \neq \tau \in \text{Rad}_\phi \cap (\bigcap_{q' \neq p \in \overline{1, 2n}} \ker_{\varphi_p})$, we have

$$\begin{aligned} \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle} &\ni [x^{-\tau - 2\varepsilon - 2\sigma_q, \mathbf{0}}, [u, x^{\tau, \mathbf{0}}]] \\ &\equiv \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\alpha, \mathbf{i}) \in \ker_{\varphi_2} \times \mathcal{J} \\ i_2 = 0, |\mathbf{i}| = \hat{k}}} a_{\alpha, \mathbf{i}} \varphi_{q'}(\tau) \varphi_q(\alpha) (\varphi_q(\alpha + \sigma_q) \varphi_{q'}(\tau + 2\sigma_q) \\ &\quad - 2\varphi_q(\sigma_q) \varphi_{q'}(\alpha + \tau + \sigma_q)) x^{\alpha, \mathbf{i}} \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}} \end{aligned} \quad (2.36)$$

(cf. (1.10)). Since $\varphi_{q'}(\tau)$ takes an infinite number of elements in $\mathbb{F}$ because $\mathbb{Z}\tau \subset \text{Rad}_\phi \cap (\cap_{q' \neq p \in \overline{1, 2n}} \ker_{\varphi_p})$, the coefficients of $\varphi_{q'}^2(\tau)$ in (2.36) show

$$\varphi_q(\alpha)\varphi_q(\alpha - \sigma_q) = \varphi_q(\beta)\varphi_q(\beta - \sigma_q) \quad \text{whenever } a_{\alpha, \mathbf{i}}a_{\beta, \mathbf{j}} \neq 0 \quad (2.37)$$

by the minimality of $\iota(u)$ and Lemma 3. Moreover, (2.37) is equivalent to

$$\varphi_q(\alpha) = \varphi_q(\beta) \text{ or } \varphi_q(\alpha + \beta - \sigma_q) = 0. \quad (2.38)$$

Assume that there exist $\alpha, \beta \in \Gamma$ such that $\varphi_q(\alpha) \neq \varphi_q(\beta)$, $\varphi_q(\alpha + \beta - \sigma_q) = 0$ and $a_{\alpha, \mathbf{i}}a_{\beta, \mathbf{j}} \neq 0$. Without loss of generality, we assume that $\varphi_q(\alpha) \neq 0$. Since $\varphi_q \not\equiv 0$ and $\varphi_{q'} \not\equiv 0$, we can choose

$$\tau \in \text{Rad}_\phi \bigcap (\bigcap_{q \neq p \in \overline{1, 2n}} \ker_{\varphi_p}) \setminus \ker_{\varphi_q}, \quad \tau' \in \text{Rad}_\phi \bigcap (\bigcap_{q' \neq p \in \overline{1, 2n}} \ker_{\varphi_p}) \setminus \ker_{\varphi_{q'}} \quad (2.39)$$

by (1.9) such that

$$\varphi_q(\tau + \alpha) \neq 0, \quad \varphi_q(\alpha)\varphi_{q'}(\tau' - \sigma_q) \neq \varphi_{q'}(\alpha)\varphi_q(\tau - \sigma_q). \quad (2.40)$$

Then we have

$$\begin{aligned} \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle} &\ni [x^{\tau + \tau' - \varepsilon - \sigma_q, \mathbf{0}}, u] \equiv \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\gamma, \mathbf{l}) \in \ker_{\varphi_2} \times \mathcal{J} \\ l_2 = 0, |\mathbf{l}| = \hat{k}}} a_{\gamma, \mathbf{l}} \epsilon_q(\varphi_q(\tau - \sigma_q)\varphi_{q'}(\gamma) \\ &\quad - \varphi_{q'}(\tau' - \sigma_q)\varphi_q(\gamma)) x^{\gamma + \tau + \tau', \mathbf{l}} \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}}. \end{aligned} \quad (2.41)$$

Since $\iota(u)$ is minimal, $\alpha + \tau + \tau' \neq -\varepsilon$ due to $\varphi_q(\alpha + \tau + \tau') = \varphi_q(\alpha + \tau) \neq 0$, and

$$a_{\alpha, \mathbf{i}} \epsilon_q(\varphi_q(\tau - \sigma_q)\varphi_{q'}(\alpha) - \varphi_{q'}(\tau' - \sigma_q)\varphi_q(\alpha)) \neq 0, \quad (2.42)$$

we have

$$a_{\beta, \mathbf{j}} \epsilon_q(\varphi_q(\tau - \sigma_q)\varphi_{q'}(\beta) - \varphi_{q'}(\tau' - \sigma_q)\varphi_q(\beta)) \neq 0. \quad (2.43)$$

But

$$\varphi_q(\alpha + \tau + \tau') \neq \varphi_q(\beta + \tau + \tau') \text{ and } \varphi_q(\alpha + \tau + \tau' + \beta + \tau + \tau' - \sigma_q) \neq 0, \quad (2.44)$$

which contradicts (2.38) if we replace u by $[x^{\tau + \tau' - \varepsilon - \sigma_q, \mathbf{0}}, u]$. Thus the first equation in (2.38) holds, and so does (2.34).

If $\varphi_q \not\equiv 0$ and $\varphi_{q'} \equiv 0$, then $\mathcal{J}_{q'} = \mathbb{N}$ by (1.7). Moreover, if $j_{q'} \neq 0$ for some $a_{\beta, \mathbf{j}} \neq 0$, choose $\tau \in \text{Rad}_\phi \cap (\cap_{q \neq p \in \overline{1, 2n}} \ker_{\varphi_p}) \setminus \ker_{\varphi_q}$ such that $\varphi_q(\beta + \tau + \sigma_q) \neq 0$. Then we have

$$\begin{aligned} \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k}-1 \rangle} &\ni [x^{\tau, \mathbf{0}}, u] \\ &\equiv \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\gamma, \mathbf{l}) \in \ker_{\varphi_2} \times \mathcal{J} \\ l_2 = 0, |\mathbf{l}| = \hat{k}}} a_{\gamma, \mathbf{l}} l_{q'} \varphi_q(\tau) x^{\gamma + \tau + \sigma_q + \varepsilon, \mathbf{l} - 1_{[q']}} \pmod{\mathcal{A}_{\langle \hat{k}-2 \rangle}}. \end{aligned} \quad (2.45)$$

Since $\varphi_q(\beta + \tau + \sigma_q + \varepsilon) = \varphi_q(\beta + \tau + \sigma_q) \neq 0$ and $j_{q'} \neq 0$ for some $a_{\beta, \mathbf{j}} \neq 0$, we have $[x^{\tau, \mathbf{0}}, u] \in (\mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k}-1 \rangle}) \setminus \mathbb{F}x^{-\varepsilon, \mathbf{0}}$, which contradicts the minimality of $\hat{k}$. Therefore $j_{q'} = 0$ whenever $a_{\beta, \mathbf{j}} \neq 0$. Thus we get

$$\begin{aligned} \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle} &\ni [x^{-\sigma_q - \varepsilon, \mathbf{1}_{[q]}}, u] \\ &\equiv \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\gamma, \mathbf{l}) \in \ker \varphi_2 \times \mathcal{J} \\ l_2=0, |\mathbf{l}|=\hat{k}}} -a_{\gamma, \mathbf{l}} \epsilon_q \varphi_q(\gamma) x^{\gamma, \mathbf{l}} \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}}, \end{aligned} \quad (2.46)$$

which implies (2.34). Therefore, we have proved

$$\varphi_q(\alpha) = \varphi_q(\beta) \quad \text{whenever } a_{\alpha, \mathbf{i}} a_{\beta, \mathbf{j}} \neq 0, \quad \text{for } q \in \overline{1, 2n}. \quad (2.47)$$

Thirdly, we want to prove that

$$\phi(\gamma, \alpha) = \phi(\gamma, \beta) \text{ for any } \gamma \in \Gamma, \text{ whenever } a_{\alpha, \mathbf{i}} a_{\beta, \mathbf{j}} \neq 0. \quad (2.48)$$

For $a_{\beta, \mathbf{j}} \neq 0$, we have

$$\begin{aligned} \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle} &\ni [x^{\beta, \mathbf{0}}, u] \\ &\equiv \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\alpha, \mathbf{i}) \in \ker \varphi_2 \times \mathcal{J} \\ i_2=0, |\mathbf{i}|=\hat{k}}} a_{\alpha, \mathbf{i}} \phi(\beta, \alpha) x^{\alpha + \beta + \varepsilon, \mathbf{i}} \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}} \end{aligned} \quad (2.49)$$

by (2.1) and (2.47). Since $\iota(u)$ is minimal and $\phi(\beta, \beta) = 0$, we have

$$\phi(\alpha, \beta) = 0 \quad \text{whenever } a_{\alpha, \mathbf{i}} a_{\beta, \mathbf{j}} \neq 0. \quad (2.50)$$

For any $\gamma \in \Gamma$, we have

$$\begin{aligned} \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle} &\ni [x^{-\gamma - \varepsilon, \mathbf{0}}, [x^{\gamma - \varepsilon, \mathbf{0}}, u]] \\ &\equiv \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\alpha, \mathbf{i}) \in \ker \varphi_2 \times \mathcal{J} \\ i_2=0, |\mathbf{i}|=\hat{k}}} a_{\alpha, \mathbf{i}} \left(\sum_{s=1}^{n-1} \sum_{t=1}^{n-1} (\varphi_{2s+1}(\gamma) \varphi_{2s+2}(\alpha) - \varphi_{2s+1}(\alpha) \varphi_{2s+2}(\gamma)) \cdot \right. \\ &\quad (\varphi_{2t+1}(\alpha + \sigma_{2s+1}) \varphi_{2t+2}(\gamma) - \varphi_{2t+1}(\gamma) \varphi_{2t+2}(\alpha + \sigma_{2s+1})) x^{\alpha + \sigma_{2s+1} + \sigma_{2t+1}, \mathbf{i}} \\ &\quad + 2 \sum_{s=1}^{n-1} (\varphi_{2s+1}(\gamma) \varphi_{2s+2}(\alpha) - \varphi_{2s+1}(\alpha) \varphi_{2s+2}(\gamma)) \varphi_2(\gamma) (\varphi_1(\alpha) + 1) x^{\alpha + \sigma_{2s+1} - \varepsilon, \mathbf{i}} \\ &\quad - 2 \sum_{s=1}^{n-1} (\varphi_{2s+1}(\gamma) \varphi_{2s+2}(\alpha) - \varphi_{2s+1}(\alpha) \varphi_{2s+2}(\gamma)) \phi(\gamma, \alpha) x^{\alpha + \sigma_{2s+1}, \mathbf{i}} \\ &\quad \left. - \phi^2(\gamma, \alpha) x^{\alpha, \mathbf{i}} + 2 \phi(\gamma, \alpha) \varphi_2(\gamma) (\varphi_1(\alpha) + 1) x^{\alpha - \varepsilon, \mathbf{i}} \right. \\ &\quad \left. - \varphi_2^2(\gamma) \varphi_1(\alpha) (\varphi_1(\alpha) + 1) x^{\alpha - 2\varepsilon, \mathbf{i}} \right) \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}}. \end{aligned} \quad (2.51)$$

For $a_{\alpha, \mathbf{i}} \neq 0$, if $\varphi_q(\alpha) = 0$ for $q \in \overline{3, 2n}$ and $\varphi_1(\alpha) + 1 = 0$, then (2.47) and (2.51) imply

$$\begin{aligned} \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle} &\ni [x^{-\gamma-\varepsilon, \mathbf{0}}, [x^{\gamma-\varepsilon, \mathbf{0}}, u]] \\ &\equiv \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\alpha, \mathbf{i}) \in \ker \varphi_2 \times \mathcal{J} \\ i_2=0, |\mathbf{i}|=\hat{k}}} -a_{\alpha, \mathbf{i}} \phi^2(\gamma, \alpha) x^{\alpha, \mathbf{i}} \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}}. \end{aligned} \quad (2.52)$$

By the minimality of $\iota(u)$ and Lemma 3, we have

$$\phi^2(\gamma, \alpha) = \phi^2(\gamma, \beta) \quad \text{whenever } a_{\alpha, \mathbf{i}} a_{\beta, \mathbf{j}} \neq 0, \quad (2.53)$$

which implies

$$\phi(\gamma, \alpha) = \phi(\gamma, \beta) \text{ or } \phi(\gamma, \alpha) = -\phi(\gamma, \beta) \quad \text{whenever } a_{\alpha, \mathbf{i}} a_{\beta, \mathbf{j}} \neq 0. \quad (2.54)$$

Assume that there exist $\alpha, \beta \in \Gamma$ such that

$$\phi(\gamma, \alpha) \neq \phi(\gamma, \beta), \quad \phi(\gamma, \alpha) = -\phi(\gamma, \beta) \text{ and } a_{\alpha, \mathbf{i}} a_{\beta, \mathbf{j}} \neq 0. \quad (2.55)$$

Then we obtain $\phi(\gamma, \alpha) = -\phi(\gamma, \beta) \neq 0$. Moreover, (2.47) and (2.50) tell us that

$$\begin{aligned} \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle} &\ni [x^{-\gamma+\alpha, \mathbf{0}}, [x^{\gamma-\varepsilon, \mathbf{0}}, u]] \\ &\equiv - \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\theta, \mathbf{i}) \in \ker \varphi_2 \times \mathcal{J} \\ i_2=0, |\mathbf{i}|=\hat{k}}} a_{\theta, \mathbf{i}} \phi(\gamma, \theta) \phi(\gamma, \theta + \alpha) x^{\theta+\alpha+\varepsilon, \mathbf{i}} \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}}. \end{aligned} \quad (2.56)$$

Since $\phi(\gamma, 2\alpha + \varepsilon) = 2\phi(\gamma, \alpha) \neq 0$, we have $2\alpha + \varepsilon \neq -\varepsilon$. But the fact that

$$a_{\alpha, \mathbf{i}} \phi(\gamma, \alpha) \phi(\gamma, 2\alpha) \neq 0, \quad a_{\beta, \mathbf{j}} \phi(\gamma, \beta) \phi(\gamma, \alpha + \beta) = 0 \quad (2.57)$$

contradicts the minimality of $\iota(u)$. Thus the first equation in (2.54) holds, and so does (2.48).

Next we assume that

$$\varphi_1(\alpha) + 1 \neq 0 \text{ or there exists } q \in \overline{3, 2n} \text{ such that } \varphi_q(\alpha) \neq 0, \text{ for } a_{\alpha, \mathbf{i}} \neq 0. \quad (2.58)$$

Claim 1. There exists some $v_1 \in \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle}$ such that

$$\begin{aligned} v_1 &\equiv \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\alpha, \mathbf{i}) \in \ker \varphi_2 \times \mathcal{J} \\ i_2=0, |\mathbf{i}|=\hat{k}}} a_{\alpha, \mathbf{i}} \left(\sum_{s=1}^{n-1} \sum_{t=1}^{n-1} (\varphi_{2s+1}(\gamma) \varphi_{2s+2}(\alpha) - \varphi_{2s+1}(\alpha) \varphi_{2s+2}(\gamma)) \cdot \right. \\ &\quad (\varphi_{2t+1}(\alpha + \sigma_{2s+1}) \varphi_{2t+2}(\gamma) - \varphi_{2t+1}(\gamma) \varphi_{2t+2}(\alpha + \sigma_{2s+1})) x^{\alpha + \sigma_{2s+1} + \sigma_{2t+1}, \mathbf{i}} \\ &\quad - 2 \sum_{s=1}^{n-1} (\varphi_{2s+1}(\gamma) \varphi_{2s+2}(\alpha) - \varphi_{2s+1}(\alpha) \varphi_{2s+2}(\gamma)) \phi(\gamma, \alpha) x^{\alpha + \sigma_{2s+1}, \mathbf{i}} \\ &\quad \left. - \phi^2(\gamma, \alpha) x^{\alpha, \mathbf{i}} \right) \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}}. \end{aligned} \quad (2.59)$$

For convenience, we denote

$$U' := \mathcal{A}_{\langle \hat{k} \rangle} \cap \left(\sum_{(\alpha, \mathbf{i}) \in \ker \varphi_2 \times \mathcal{J}; i_2=0} \mathbb{F}x^{\alpha, \mathbf{i}} \right), \quad (2.60)$$

$$W_1 := \mathcal{A}_{\langle \hat{k}-1 \rangle} \cap \left(\sum_{(\alpha, \mathbf{i}) \in \ker \varphi_2 \times \mathcal{J}; i_2=0} \mathbb{F}x^{\alpha, \mathbf{i}} \right), \quad (2.61)$$

$$W_2 := \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle} = \mathcal{I} \cap U'. \quad (2.62)$$

Notice that W_1 and W_2 are two subspaces of U' . If $\mathcal{J}_2 = \mathbb{N}$, $\text{ad}_{x^{0,1[2]}}$ is a linear transformation on $\mathcal{A}$ which preserves U' , W_1 and W_2 . For any $a_{\alpha, \mathbf{i}} \neq 0$, we have

$$[x^{0,1[2]}, x^{\alpha, \mathbf{i}}] \equiv -(\varphi_1(\alpha) + 1)x^{\alpha, \mathbf{i}} \pmod{W_1}. \quad (2.63)$$

Thus, acting on U'/W_1 , $\text{ad}_{x^{0,1[2]}}$ has eigenvectors $x^{\alpha+\sigma_{2s+1}+\sigma_{2t+1}, \mathbf{i}} + W_1$, $x^{\alpha+\sigma_{2s+1}, \mathbf{i}} + W_1$ and $x^{\alpha, \mathbf{i}} + W_1$ with eigenvalue $-\varphi_1(\alpha) - 1$. Moreover, it has eigenvectors $x^{\alpha+\sigma_{2s+1}-\varepsilon, \mathbf{i}} + W_1$ and $x^{\alpha-\varepsilon, \mathbf{i}} + W_1$ with eigenvalue $-\varphi_1(\alpha)$. Furthermore, $x^{\alpha-2\varepsilon, \mathbf{i}} + W_1$ is an eigenvector with eigenvalue $-\varphi_1(\alpha) + 1$. Repeatedly applying $\text{ad}_{x^{0,1[2]}}$ on $(W_2 + W_1)/W_1$, *Claim 1* holds by Lemma 3, (2.47) and (2.51).

If $\mathcal{J}_2 = \{0\}$, we have $\varphi_2 \not\equiv 0$ by (1.7). For any $a_{\alpha, \mathbf{i}} \neq 0$, if $\varphi_1(\alpha) + 1 = 0$, *Claim 1* follows naturally from (2.51). Suppose that $\varphi_1(\alpha) + 1 \neq 0$. Picking $\tau \in \text{Rad}_\phi \cap (\cap_{2 \neq p \in \overline{1, 2n}} \ker \varphi_p) \setminus \ker \varphi_2$, $\text{ad}_{x^{-\tau, \mathbf{0}}} \text{ad}_{x^{\tau, \mathbf{0}}}$ is a linear transformation on $\mathcal{A}$ which preserves U' , W_1 and W_2 . Since for any $a_{\alpha, \mathbf{i}} \neq 0$, we have

$$[x^{-\tau, \mathbf{0}}, [x^{\tau, \mathbf{0}}, x^{\alpha, \mathbf{i}}]] \equiv g(\alpha)x^{\alpha, \mathbf{i}} \pmod{W_1}, \quad (2.64)$$

where $g(\alpha) = -\varphi_2^2(\tau)(\varphi_1(\alpha) + 1)(\varphi_1(\alpha) + 2)$. Acting on U'/W_1 , $\text{ad}_{x^{-\tau, \mathbf{0}}} \text{ad}_{x^{\tau, \mathbf{0}}}$ has eigenvectors $x^{\alpha+\sigma_{2s+1}+\sigma_{2t+1}, \mathbf{i}} + W_1$, $x^{\alpha+\sigma_{2s+1}, \mathbf{i}} + W_1$ and $x^{\alpha, \mathbf{i}} + W_1$ with eigenvalue $g(\alpha)$. Moreover, it has eigenvectors $x^{\alpha+\sigma_{2s+1}-\varepsilon, \mathbf{i}} + W_1$ and $x^{\alpha-\varepsilon, \mathbf{i}} + W_1$ with eigenvalue $g(\alpha - \varepsilon)$. Furthermore, $x^{\alpha-2\varepsilon, \mathbf{i}} + W_1$ is an eigenvector with eigenvalue $g(\alpha - 2\varepsilon)$. First, we have $g(\alpha) \neq g(\alpha - \varepsilon)$. If $2\varphi_1(\alpha) + 1 \neq 0$, we further get $g(\alpha) \neq g(\alpha - 2\varepsilon)$. Thus, repeatedly applying $\text{ad}_{x^{-\tau, \mathbf{0}}} \text{ad}_{x^{\tau, \mathbf{0}}}$ on $(W_2 + W_1)/W_1$, *Claim 1* holds by Lemma 3, (2.47) and (2.51). Otherwise, $2\varphi_1(\alpha) + 1 = 0$, we have $g(\alpha) = g(\alpha - 2\varepsilon)$. By Lemma 3, (2.47) and (2.51) we obtain some $v'_1 \in \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle}$ that

$$\begin{aligned} v'_1 &\equiv \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\alpha, \mathbf{i}) \in \ker \varphi_2 \times \mathcal{J} \\ i_2=0, |\mathbf{i}|=\hat{k}}} a_{\alpha, \mathbf{i}} \left(\sum_{s=1}^{n-1} \sum_{t=1}^{n-1} (\varphi_{2s+1}(\gamma)\varphi_{2s+2}(\alpha) - \varphi_{2s+1}(\alpha)\varphi_{2s+2}(\gamma)) \cdot \right. \\ &\quad (\varphi_{2t+1}(\alpha + \sigma_{2s+1})\varphi_{2t+2}(\gamma) - \varphi_{2t+1}(\gamma)\varphi_{2t+2}(\alpha + \sigma_{2s+1})) x^{\alpha+\sigma_{2s+1}+\sigma_{2t+1}, \mathbf{i}} \\ &\quad - 2 \sum_{s=1}^{n-1} (\varphi_{2s+1}(\gamma)\varphi_{2s+2}(\alpha) - \varphi_{2s+1}(\alpha)\varphi_{2s+2}(\gamma)) \phi(\gamma, \alpha) x^{\alpha+\sigma_{2s+1}, \mathbf{i}} \\ &\quad \left. - \phi^2(\gamma, \alpha) x^{\alpha, \mathbf{i}} - \varphi_2^2(\gamma)\varphi_1(\alpha)(\varphi_1(\alpha) + 1)x^{\alpha-2\varepsilon, \mathbf{i}} \right) \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}}. \quad (2.65) \end{aligned}$$

Since $2\varphi_1(\alpha) + 1 = 0$, we have $\varphi_1(\alpha) \neq 0$. Pick $\tau \in \text{Rad}_\phi \cap (\cap_{2 \neq p \in \overline{1, 2n}} \ker_{\varphi_p}) \setminus \ker_{\varphi_2}$ again, then we have

$$\begin{aligned} \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle} &\ni [x^{-\tau, \mathbf{0}}, [x^{\tau-2\varepsilon, \mathbf{0}}, u]] \equiv \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\alpha, \mathbf{i}) \in \ker_{\varphi_2} \times \mathcal{J} \\ i_2=0, |\mathbf{i}|=\hat{k}}} -a_{\alpha, \mathbf{i}} \varphi_2^2(\tau) \varphi_1(\alpha) \cdot \\ &(\varphi_1(\alpha) + 1)x^{\alpha-2\varepsilon, \mathbf{i}} \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}}. \end{aligned} \quad (2.66)$$

Therefore, *Claim 1* holds by substituting (2.65) and (2.66) in

$$v'_1 - \frac{\varphi_2^2(\gamma)}{\varphi_2^2(\tau)} [x^{-\tau, \mathbf{0}}, [x^{\tau-2\varepsilon, \mathbf{0}}, u]]. \quad (2.67)$$

Claim 2. For any fixed r (where $r \in \overline{1, n-1}$), there exists some $v_2 \in \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle}$ such that

$$\begin{aligned} v_2 &\equiv \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\alpha, \mathbf{i}) \in \ker_{\varphi_2} \times \mathcal{J} \\ i_2=0, |\mathbf{i}|=\hat{k}}} a_{\alpha, \mathbf{i}} \left(\sum_{\substack{s=1 \\ s \neq r}}^{n-1} \sum_{\substack{t=1 \\ t \neq r}}^{n-1} (\varphi_{2s+1}(\gamma) \varphi_{2s+2}(\alpha) - \varphi_{2s+1}(\alpha) \varphi_{2s+2}(\gamma)) \cdot \right. \\ &(\varphi_{2t+1}(\alpha + \sigma_{2s+1}) \varphi_{2t+2}(\gamma) - \varphi_{2t+1}(\gamma) \varphi_{2t+2}(\alpha + \sigma_{2s+1})) x^{\alpha + \sigma_{2s+1} + \sigma_{2t+1}, \mathbf{i}} \\ &- 2 \sum_{\substack{s=1 \\ s \neq r}}^{n-1} (\varphi_{2s+1}(\gamma) \varphi_{2s+2}(\alpha) - \varphi_{2s+1}(\alpha) \varphi_{2s+2}(\gamma)) \phi(\gamma, \alpha) x^{\alpha + \sigma_{2s+1}, \mathbf{i}} \\ &\left. - \phi^2(\gamma, \alpha) x^{\alpha, \mathbf{i}} \right) \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}}. \end{aligned} \quad (2.68)$$

If $\sigma_{2r+1} = 0$, we get $\varphi_{2r+1} \equiv \varphi_{2r+2} \equiv 0$ by (1.10). Therefore *Claim 2* follows naturally from (2.59). We assume that $\sigma_{2r+1} \neq 0$. Then $\varphi_{2r+1}(\sigma_{2r+1}) \neq 0$ or $\varphi_{2r+2}(\sigma_{2r+1}) \neq 0$ by (1.8) and (1.10). Choose $q \in \{2r+1, 2r+2\}$ such that $\varphi_q(\sigma_{2r+1}) \neq 0$, then $\varphi_q \not\equiv 0$ and $\sigma_q = \sigma_{2r+1}$.

If $\varphi_{q'} \equiv 0$, we get $\mathcal{J}_{q'} = \mathbb{N}$ by (1.7). Observe that $\text{ad}_{x^{-\sigma_q - \varepsilon, 1_{[q']}}}$ is a linear transformation on $\mathcal{A}$ which preserves U' , W_1 and W_2 . For any $a_{\alpha, \mathbf{i}} \neq 0$, since $i_{q'} = 0$ by (2.45), we have

$$[x^{-\sigma_q - \varepsilon, 1_{[q']}}, x^{\alpha, \mathbf{i}}] \equiv -\epsilon_q \varphi_q(\alpha) x^{\alpha, \mathbf{i}} \pmod{W_1}. \quad (2.69)$$

Repeatedly applying $\text{ad}_{x^{-\sigma_q - \varepsilon, 1_{[q']}}}$ on $(W_2 + W_1)/W_1$, *Claim 2* holds by Lemma 3, (2.47), (2.59) and (2.69).

If $\varphi_{q'} \not\equiv 0$, pick $\tau \in \text{Rad}_\phi \cap (\cap_{q' \neq p \in \overline{1, 2n}} \ker_{\varphi_p}) \setminus \ker_{\varphi_{q'}}$. Then $\text{ad}_{x^{-\tau - 2\varepsilon - 2\sigma_q, \mathbf{0}}} \text{ad}_{x^{\tau, \mathbf{0}}}$ is a linear transformation on $\mathcal{A}$ which preserves U' , W_1 and W_2 . For any $a_{\alpha, \mathbf{i}} \neq 0$, we have

$$[x^{-\tau - 2\varepsilon - 2\sigma_q, \mathbf{0}}, [x^{\tau, \mathbf{0}}, x^{\alpha, \mathbf{i}}]] \equiv h(\alpha) x^{\alpha, \mathbf{i}} \pmod{W_1}, \quad (2.70)$$

where $h(\alpha) = -\varphi_{q'}(\tau)\varphi_q(\alpha)(\varphi_q(\alpha)\varphi_{q'}(\tau + 2\sigma_q) - \varphi_q(\sigma_q)\varphi_{q'}(\tau + 2\alpha))$. Acting on U'/W_1 , $\text{ad}_{x^{-\tau-2\varepsilon-2\sigma_q}, \mathbf{0}} \text{ad}_{x^{\tau, \mathbf{0}}}$ has eigenvectors $x^{\alpha, \mathbf{i}} + W_1$, $x^{\alpha+\sigma_{2s+1}+\sigma_{2t+1}, \mathbf{i}} + W_1$ and $x^{\alpha+\sigma_{2s+1}, \mathbf{i}} + W_1$ (for $s, t \neq r$) with eigenvalue $h(\alpha)$. Moreover, it has eigenvectors $x^{\alpha+\sigma_{2r+1}, \mathbf{i}} + W_1$ and $x^{\alpha+\sigma_{2s+1}+\sigma_{2r+1}, \mathbf{i}} + W_1$ (for $s \neq r$) with eigenvalue $h(\alpha + \sigma_q)$. Furthermore, $x^{\alpha+2\sigma_{2r+1}, \mathbf{i}} + W_1$ is an eigenvector with eigenvalue $h(\alpha + 2\sigma_q)$.

For any $a_{\alpha, \mathbf{i}} \neq 0$, if $\varphi_q(\alpha) = \varphi_{q'}(\alpha) = 0$, *Claim 2* follows from (2.59). Assume that $\varphi_q(\alpha) \neq 0$ or $\varphi_{q'}(\alpha) \neq 0$. Restricting the selection of τ so that

$$\varphi_q(\alpha)\varphi_{q'}(\tau + \sigma_q) - \varphi_q(\sigma_q)\varphi_{q'}(\alpha) \neq 0, \quad (2.71)$$

we have $h(\alpha) \neq h(\alpha + \sigma_q)$. If $\varphi_q(2\alpha + \sigma_q) \neq 0$ or $\varphi_{q'}(2\alpha + \sigma_q) \neq 0$, we can further restrict the selection of τ so that

$$\varphi_q(2\alpha + \sigma_q)\varphi_{q'}(\tau + \sigma_q) - \varphi_q(\sigma_q)\varphi_{q'}(2\alpha + \sigma_q) \neq 0. \quad (2.72)$$

Therefore we have $h(\alpha) \neq h(\alpha + 2\sigma_q)$. Repeatedly applying $\text{ad}_{x^{-\tau-2\varepsilon-2\sigma_q}, \mathbf{0}} \text{ad}_{x^{\tau, \mathbf{0}}}$ on $(W_2 + W_1)/W_1$, *Claim 2* follows from Lemma 3, (2.47) and (2.59). Otherwise, $\varphi_q(2\alpha + \sigma_q) = \varphi_{q'}(2\alpha + \sigma_q) = 0$, we get $h(\alpha) = h(\alpha + 2\sigma_q)$. By Lemma 3, (2.47) and (2.59), we obtain some $v'_2 \in \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle}$ such that

$$\begin{aligned} v'_2 \equiv & \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\alpha, \mathbf{i}) \in \ker \varphi_2 \times \mathcal{J} \\ i_2=0, |\mathbf{i}|=\hat{k}}} a_{\alpha, \mathbf{i}} \left(\sum_{\substack{s=1 \\ s \neq r}}^{n-1} \sum_{\substack{t=1 \\ t \neq r}}^{n-1} (\varphi_{2s+1}(\gamma)\varphi_{2s+2}(\alpha) - \varphi_{2s+1}(\alpha)\varphi_{2s+2}(\gamma)) \cdot \right. \\ & (\varphi_{2t+1}(\alpha + \sigma_{2s+1})\varphi_{2t+2}(\gamma) - \varphi_{2t+1}(\gamma)\varphi_{2t+2}(\alpha + \sigma_{2s+1})) x^{\alpha+\sigma_{2s+1}+\sigma_{2t+1}, \mathbf{i}} \\ & - 2 \sum_{\substack{s=1 \\ s \neq r}}^{n-1} (\varphi_{2s+1}(\gamma)\varphi_{2s+2}(\alpha) - \varphi_{2s+1}(\alpha)\varphi_{2s+2}(\gamma)) \phi(\gamma, \alpha) x^{\alpha+\sigma_{2s+1}, \mathbf{i}} \\ & + (\varphi_{2r+1}(\gamma)\varphi_{2r+2}(\alpha) - \varphi_{2r+1}(\alpha)\varphi_{2r+2}(\gamma))^2 x^{\alpha+2\sigma_{2r+1}, \mathbf{i}} \\ & \left. - \phi^2(\gamma, \alpha) x^{\alpha, \mathbf{i}} \right) \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}}. \end{aligned} \quad (2.73)$$

Since $\varphi_q(2\alpha + \sigma_q) = 0$ and $\varphi_q(\sigma_q) \neq 0$, we have $\varphi_q(\alpha) \neq 0$ and $\varphi_q(\alpha + \sigma_q) \neq 0$. Picking $\tau' \in \text{Rad}_\phi \cap (\cap_{q' \neq p \in \overline{1, 2n}} \ker \varphi_p) \setminus \ker \varphi_{q'}$, we have

$$\begin{aligned} \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle} \ni & [x^{-\tau'-\varepsilon, \mathbf{0}}, [x^{\tau'-\varepsilon, \mathbf{0}}, u]] \equiv \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\alpha, \mathbf{i}) \in \ker \varphi_2 \times \mathcal{J} \\ i_2=0, |\mathbf{i}|=\hat{k}}} -a_{\alpha, \mathbf{i}} \varphi_{q'}^2(\tau') \varphi_q(\alpha) \cdot \\ & \varphi_q(\alpha + \sigma_q) x^{\alpha+2\sigma_q, \mathbf{i}} \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}}. \end{aligned} \quad (2.74)$$

Thus, *Claim 2* holds by substituting (2.73) and (2.74) in

$$v'_2 + \frac{(\varphi_{2r+1}(\gamma)\varphi_{2r+2}(\alpha) - \varphi_{2r+1}(\alpha)\varphi_{2r+2}(\gamma))^2}{\varphi_{q'}^2(\tau')\varphi_q(\alpha)\varphi_q(\alpha + \sigma_q)} [x^{-\tau'-\varepsilon, \mathbf{0}}, [x^{\tau'-\varepsilon, \mathbf{0}}, u]]. \quad (2.75)$$

Claim 3. There exists some $v_n \in \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle}$ such that

$$v_n \equiv \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\alpha, \mathbf{i}) \in \ker \varphi_2 \times \mathcal{J} \\ i_2=0, |\mathbf{i}|=\hat{k}}} -a_{\alpha, \mathbf{i}} \phi^2(\gamma, \alpha) x^{\alpha, \mathbf{i}} \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}}. \quad (2.76)$$

For all $r \in \overline{1, n-1}$, repeat the procedure of the proof of *Claim 2*. It is clear that *Claim 3* holds.

Hence, by the minimality of $\iota(u)$, *Claim 3* and Lemma 3, we get

$$\phi^2(\gamma, \alpha) = \phi^2(\gamma, \beta) \quad \text{whenever } a_{\alpha, \mathbf{i}} a_{\beta, \mathbf{j}} \neq 0, \quad (2.77)$$

namely,

$$\phi(\gamma, \alpha) = \phi(\gamma, \beta) \text{ or } \phi(\gamma, \alpha) = -\phi(\gamma, \beta) \quad \text{whenever } a_{\alpha, \mathbf{i}} a_{\beta, \mathbf{j}} \neq 0. \quad (2.78)$$

Assume that there exist $\alpha, \beta \in \Gamma$ such that

$$\phi(\gamma, \alpha) \neq \phi(\gamma, \beta), \quad \phi(\gamma, \alpha) = -\phi(\gamma, \beta) \text{ and } a_{\alpha, \mathbf{i}} a_{\beta, \mathbf{j}} \neq 0. \quad (2.79)$$

Then we obtain $\phi(\gamma, \alpha) = -\phi(\gamma, \beta) \neq 0$.

For any $a_{\alpha, \mathbf{i}} \neq 0$, if there exists $q \in \overline{3, 2n}$ such that $\varphi_q(\alpha) \neq 0$ and $\varphi_{q'} \not\equiv 0$. Picking $\tau' \in \text{Rad}_\phi \cap (\cap_{q' \neq p \in \overline{1, 2n}} \ker \varphi_p) \setminus \ker \varphi_{q'}$, we have

$$\begin{aligned} \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle} &\ni [x^{\alpha + \tau', \mathbf{0}}, u] \\ &\equiv \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\theta, \mathbf{l}) \in \ker \varphi_2 \times \mathcal{J} \\ l_2=0, |\mathbf{l}|=\hat{k}}} a_{\theta, \mathbf{l}} \epsilon_{q'} \varphi_{q'}(\tau') \varphi_q(\theta) x^{\theta + \alpha + \tau' + \sigma_q + \varepsilon, \mathbf{l}} \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}}. \end{aligned} \quad (2.80)$$

Since $2\alpha + \tau' + \sigma_q + \varepsilon \neq -\varepsilon$ due to $\phi(\gamma, 2\alpha + \tau' + \sigma_q + \varepsilon) = 2\phi(\gamma, \alpha) \neq 0$, and $a_{\alpha, \mathbf{i}} \epsilon_{q'} \varphi_{q'}(\tau') \varphi_q(\alpha) \neq 0$, we have $[x^{\alpha + \tau', \mathbf{0}}, u] \in (\mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle}) \setminus \mathbb{F} x^{-\varepsilon, \mathbf{0}}$. Then the minimality of $\iota(u)$ implies that $\iota([x^{\alpha + \tau', \mathbf{0}}, u]) = \iota(u)$. But we have

$$\phi(\gamma, 2\alpha + \tau' + \sigma_q + \varepsilon) \neq \phi(\gamma, \beta + \alpha + \tau' + \sigma_q + \varepsilon) \quad (2.81)$$

and

$$\phi(\gamma, 2\alpha + \tau' + \sigma_q + \varepsilon) + \phi(\gamma, \beta + \alpha + \tau' + \sigma_q + \varepsilon) = 2\phi(\gamma, \alpha) \neq 0, \quad (2.82)$$

which contradicts (2.78) if we replace u by $[x^{\alpha + \tau', \mathbf{0}}, u]$. Thus the first equation in (2.78) holds, and so does (2.48).

For any $a_{\alpha, \mathbf{i}} \neq 0$, if either $\varphi_p(\alpha) = 0$ or $\varphi_{p'} \equiv 0$ for $p \in \overline{3, 2n}$, and there does exist $q \in \overline{3, 2n}$ such that $\varphi_q(\alpha) \neq 0$. Fixing p , if $\varphi_p(\alpha) \neq 0$, we have $\varphi_{p'} \equiv 0$.

Then $\mathcal{J}_{p'} = \mathbb{N}$ by (1.7). By (2.45), $a_{\beta, \mathbf{j}} \neq 0$ implies that $j_{p'} = 0$. In the following, we would like to rewrite u' in (2.18) as

$$u' = \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\beta, \mathbf{l}) \in \ker \varphi_2 \times \mathcal{J} \\ l_2=0, |\mathbf{l}|=\hat{k}-1}} c_{\beta, \mathbf{l}} x^{\beta, \mathbf{l}} + u'', \quad (2.83)$$

where $c_{\beta, \mathbf{l}} \in \mathbb{F}$ and $u'' \in \mathcal{A}_{\langle \hat{k}-2 \rangle} + \mathbb{F}x^{-\varepsilon, \mathbf{0}}$. Thus we get

$$\begin{aligned} \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k}-1 \rangle} &\ni [x^{\alpha, \mathbf{0}}, u] \\ &\equiv \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\theta', \mathbf{l}') \in \ker \varphi_2 \times \mathcal{J} \\ l'_2=0, |\mathbf{l}'|=\hat{k}-1}} c_{\theta', \mathbf{l}'} \phi(\alpha, \theta') x^{\theta' + \alpha + \varepsilon, \mathbf{l}'} \pmod{\mathcal{A}_{\langle \hat{k}-2 \rangle}}. \end{aligned} \quad (2.84)$$

For any $c_{\theta', \mathbf{l}'} \neq 0$, if $\phi(\alpha, \theta') \neq 0$, we have $\theta' + \alpha + \varepsilon \neq -\varepsilon$ due to $\phi(\alpha, \theta' + \alpha + \varepsilon) \neq 0$. Then $[x^{\alpha, \mathbf{0}}, u] \in (\mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k}-1 \rangle}) \setminus \mathbb{F}x^{-\varepsilon, \mathbf{0}}$, which contradicts the minimality of $\hat{k}$ in (2.17). Therefore $\phi(\alpha, \theta') = 0$ whenever $c_{\theta', \mathbf{l}'} \neq 0$. Hence we have

$$\begin{aligned} \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle} &\ni [x^{\alpha, \mathbf{1}_{[q']}}, u] \\ &\equiv \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\theta, \mathbf{l}) \in \ker \varphi_2 \times \mathcal{J} \\ l_2=0, |\mathbf{l}|=\hat{k}}} a_{\theta, \mathbf{l}} \epsilon_{q'} \varphi_q(\theta) x^{\theta + \alpha + \sigma_q + \varepsilon, \mathbf{l}} \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}}. \end{aligned} \quad (2.85)$$

Since $2\alpha + \sigma_q + \varepsilon \neq -\varepsilon$ due to $\phi(\gamma, 2\alpha + \sigma_q + \varepsilon) \neq 0$, and $a_{\alpha, \mathbf{i}} \epsilon_{q'} \varphi_q(\alpha) \neq 0$, we have $[x^{\alpha, \mathbf{1}_{[q']}}, u] \in (\mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle}) \setminus \mathbb{F}x^{-\varepsilon, \mathbf{0}}$. Then the minimality of $\iota(u)$ implies that $\iota([x^{\alpha, \mathbf{1}_{[q']}}, u]) = \iota(u)$. But we have

$$\phi(\gamma, 2\alpha + \sigma_q + \varepsilon) \neq \phi(\gamma, \beta + \alpha + \sigma_q + \varepsilon) \quad (2.86)$$

and

$$\phi(\gamma, 2\alpha + \sigma_q + \varepsilon) + \phi(\gamma, \beta + \alpha + \sigma_q + \varepsilon) = 2\phi(\gamma, \alpha) \neq 0, \quad (2.87)$$

which contradicts (2.78) if we replace u by $[x^{\alpha, \mathbf{1}_{[q']}}, u]$. Therefore the first equation in (2.78) holds, and so does (2.48).

For any $a_{\alpha, \mathbf{i}} \neq 0$, if $\varphi_q(\alpha) = 0$ for all $q \in \overline{3, 2n}$, we therefore have $\varphi_1(\alpha) + 1 \neq 0$ by (2.58). Moreover, if $\varphi_2 \not\equiv 0$, picking $\tau \in \text{Rad}_\phi \cap (\cap_{2 \neq p \in \overline{1, 2n}} \ker \varphi_p) \setminus \ker \varphi_2$, we have

$$\begin{aligned} \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle} &\ni [x^{-\tau, \mathbf{0}}, [x^{\alpha + \tau, \mathbf{0}}, u]] \\ &\equiv \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\theta, \mathbf{l}) \in \ker \varphi_2 \times \mathcal{J} \\ l_2=0, |\mathbf{l}|=\hat{k}}} -2a_{\theta, \mathbf{l}} \varphi_2^2(\tau) (\varphi_1(\alpha) + 1)^2 x^{\theta + \alpha, \mathbf{l}} \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}}. \end{aligned} \quad (2.88)$$

Since $2\alpha \neq -\varepsilon$ due to $\phi(\gamma, 2\alpha) = 2\phi(\gamma, \alpha) \neq 0$, and $a_{\alpha, \mathbf{i}}\varphi_2^2(\tau)(\varphi_1(\alpha) + 1)^2 \neq 0$, we have $[x^{-\tau, \mathbf{0}}, [x^{\alpha+\tau, \mathbf{0}}, u]] \in (\mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle}) \setminus \mathbb{F}x^{-\varepsilon, \mathbf{0}}$. Then the minimality of $\iota(u)$ implies that $\iota([x^{-\tau, \mathbf{0}}, [x^{\alpha+\tau, \mathbf{0}}, u]]) = \iota(u)$. But we have

$$\phi(\gamma, 2\alpha) \neq \phi(\gamma, \beta + \alpha) \text{ and } \phi(\gamma, 2\alpha) + \phi(\gamma, \beta + \alpha) = 2\phi(\gamma, \alpha) \neq 0, \quad (2.89)$$

which contradicts (2.78) if we replace u by $[x^{-\tau, \mathbf{0}}, [x^{\alpha+\tau, \mathbf{0}}, u]]$. Thus the first equation in (2.78) holds, and so does (2.48). Suppose that $\varphi_2 \equiv 0$, then $\mathcal{J}_2 = \mathbb{N}$ by (1.7). By (2.50) and (2.83) we have

$$\begin{aligned} & \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k}-1 \rangle} \ni [x^{\alpha, \mathbf{0}}, u] \\ \equiv & \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\theta', \mathbf{l}') \in \ker \varphi_2 \times \mathcal{J} \\ l'_2=0, |\mathbf{l}'|=\hat{k}-1}} c_{\theta', \mathbf{l}'} \phi(\alpha, \theta') x^{\theta'+\alpha+\varepsilon, \mathbf{l}'} \pmod{\mathcal{A}_{\langle \hat{k}-2 \rangle}}. \end{aligned} \quad (2.90)$$

If $\phi(\alpha, \theta') \neq 0$ for some $c_{\theta', \mathbf{l}'} \neq 0$, we have $\theta' + \alpha + \varepsilon \neq -\varepsilon$ due to $\phi(\alpha, \theta' + \alpha + \varepsilon) \neq 0$. Then we obtain that $[x^{\alpha, \mathbf{0}}, u] \in (\mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k}-1 \rangle}) \setminus \mathbb{F}x^{-\varepsilon, \mathbf{0}}$, which contradicts the minimality of $\hat{k}$ in (2.17). Therefore $\phi(\alpha, \theta') = 0$ whenever $c_{\theta', \mathbf{l}'} \neq 0$. Similarly, we have $\phi(\alpha, \theta) = 0$ for any nonzero terms $c_{\theta, \mathbf{l}} x^{\theta, \mathbf{l}}$ of u with $|\mathbf{l}| \leq \hat{k} - 2$. Thus we get

$$\begin{aligned} & \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle} \ni [x^{\alpha, \mathbf{1}^{[2]}}, u] \\ \equiv & \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\theta, \mathbf{l}) \in \ker \varphi_2 \times \mathcal{J} \\ l_2=0, |\mathbf{l}|=\hat{k}}} -a_{\theta, \mathbf{l}}(\varphi_1(\theta) + 1) x^{\theta+\alpha, \mathbf{l}} \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}}. \end{aligned} \quad (2.91)$$

Since $2\alpha \neq -\varepsilon$ due to $\phi(\gamma, 2\alpha) = 2\phi(\gamma, \alpha) \neq 0$, and $a_{\alpha, \mathbf{i}}(\varphi_1(\alpha) + 1) \neq 0$, we have $[x^{\alpha, \mathbf{1}^{[2]}}, u] \in (\mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle}) \setminus \mathbb{F}x^{-\varepsilon, \mathbf{0}}$. Then the minimality of $\iota(u)$ implies that $\iota([x^{\alpha, \mathbf{1}^{[2]}}, u]) = \iota(u)$. But we have

$$\phi(\gamma, 2\alpha) \neq \phi(\gamma, \beta + \alpha) \text{ and } \phi(\gamma, 2\alpha) + \phi(\gamma, \beta + \alpha) = 2\phi(\gamma, \alpha) \neq 0, \quad (2.92)$$

which contradicts (2.78) if we replace u by $[x^{\alpha, \mathbf{1}^{[2]}}, u]$. Thus the first equation in (2.78) holds, and so does (2.48). Therefore we have proved that

$$\phi(\gamma, \alpha) = \phi(\gamma, \beta) \text{ for any } \gamma \in \Gamma, \quad \text{whenever } a_{\alpha, \mathbf{i}} a_{\beta, \mathbf{j}} \neq 0. \quad (2.93)$$

Hence by (1.8), (2.47) and (2.93), we get

$$\alpha = \beta \quad \text{whenever } a_{\alpha, \mathbf{i}} a_{\beta, \mathbf{j}} \neq 0. \quad (2.94)$$

Step 2. $1 \in \mathcal{I}$.

Let $a_{\alpha,i} \neq 0$ be fixed. We will give the proof in two cases.

Case 1. $\hat{k} = 0$.

In this case, $x^{\alpha,0} \in \mathcal{I}_{[0]}$ for some $\alpha \in \ker_{\varphi_2} \setminus \{-\varepsilon\}$. Assume that $\alpha \neq 0$. Since $\alpha \neq -\varepsilon$, we have $\varphi_1(\alpha) + 1 \neq 0$ or $\varphi_q(\alpha) \neq 0$ for some $q \in \overline{3, 2n}$, or $\alpha \notin \text{Rad}_\phi$.

Subcase 1.1. $\varphi_1(\alpha) + 1 \neq 0$.

If $\varphi_2 \not\equiv 0$, picking $\tau \in \text{Rad}_\phi \cap (\cap_{2 \neq p \in \overline{1, 2n}} \ker_{\varphi_p}) \setminus \ker_{\varphi_2}$, we have

$$[[x^{\alpha,0}, x^{-\alpha+\tau,0}], x^{-\tau,0}] = -2\varphi_2^2(\tau)(\varphi_1(\alpha) + 1)x^{0,0} \in \mathcal{I}. \quad (2.95)$$

Thus we get $1 \in \mathcal{I}$. Otherwise $\varphi_2 \equiv 0$, we have $\mathcal{J}_2 = \mathbb{N}$. Then we get

$$[x^{\alpha,0}, x^{-\alpha,1[2]}] = (\varphi_1(\alpha) + 1)x^{0,0} \in \mathcal{I}. \quad (2.96)$$

So we obtain $1 \in \mathcal{I}$.

Subcase 1.2. $\varphi_1(\alpha) + 1 = 0$ and $\varphi_q(\alpha) \neq 0$ for some $q \in \overline{3, 2n}$.

If $\varphi_{q'} \not\equiv 0$, picking $\tau' \in \text{Rad}_\phi \cap (\cap_{q' \neq p \in \overline{1, 2n}} \ker_{\varphi_p}) \setminus \ker_{\varphi_{q'}}$, we have

$$[x^{\alpha,0}, x^{\tau',0}] = \epsilon_q \varphi_{q'}(\tau') \varphi_q(\alpha) x^{\alpha+\tau'+\sigma_q+\varepsilon,0} \in \mathcal{I}. \quad (2.97)$$

Then $x^{\alpha+\tau'+\sigma_q+\varepsilon,0} \in \mathcal{I}$. Since $\varphi_1(\alpha+\tau'+\sigma_q+\varepsilon)+1 \neq 0$ and $\varphi_2(\alpha+\tau'+\sigma_q+\varepsilon) = 0$, we go back to *Subcase 1.1* with $x^{\alpha,0}$ replaced by $x^{\alpha+\tau'+\sigma_q+\varepsilon,0}$. Assume $\varphi_{q'} \equiv 0$, then $\mathcal{J}_{q'} = \mathbb{N}$. So we have

$$[x^{\alpha,0}, x^{0,1[q]}] = \epsilon_q \varphi_q(\alpha) x^{\alpha+\sigma_q+\varepsilon,0} \in \mathcal{I}, \quad (2.98)$$

which implies $x^{\alpha+\sigma_q+\varepsilon,0} \in \mathcal{I}$. Since $\varphi_1(\alpha+\sigma_q+\varepsilon)+1 \neq 0$ and $\varphi_2(\alpha+\sigma_q+\varepsilon) = 0$, we go back to *Subcase 1.1* with $x^{\alpha,0}$ replaced by $x^{\alpha+\sigma_q+\varepsilon,0}$.

Subcase 1.3. $\varphi_1(\alpha) + 1 = 0$ and $\varphi_q(\alpha) = 0$ for all $q \in \overline{3, 2n}$.

Since $\alpha \neq -\varepsilon$, we have $\alpha \notin \text{Rad}_\phi$. Namely, there exists $\beta \in \Gamma$ such that $\phi(\alpha, \beta) \neq 0$. Fixing any such β , we have

$$[x^{\alpha,0}, x^{\beta-\alpha-\varepsilon,0}] = \phi(\alpha, \beta) x^{\beta,0} \in \mathcal{I}, \quad (2.99)$$

which implies $x^{\beta,0} \in \mathcal{I}$. If $\varphi_1(\beta) + 1 \neq 0$ and $\varphi_2(\beta) = 0$, we go back to *Subcase 1.1* with $x^{\alpha,0}$ replaced by $x^{\beta,0}$. If $\varphi_1(\beta) + 1 \neq 0$ and $\varphi_2(\beta) \neq 0$, we have

$$[x^{\beta,0}, x^{-\beta,0}] = -2\varphi_2(\beta) x^{0,0} \in \mathcal{I}. \quad (2.100)$$

Then we get $1 \in \mathcal{I}$. If $\varphi_1(\beta) + 1 = 0$, we have $\varphi_1(\beta+\varepsilon)+1 \neq 0$ and $\phi(\alpha, \beta+\varepsilon) \neq 0$. We go back to (2.99) and (2.100) with β replaced by $\beta + \varepsilon$.

Case 2. $\hat{k} > 0$.

We have obtained that

$$u = \sum_{\substack{\mathbf{i} \in \mathcal{J} \\ i_2=0, |\mathbf{i}|=\hat{k}}} a_{\alpha, \mathbf{i}} x^{\alpha, \mathbf{i}} + u' \in (\mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle}) \setminus \mathbb{F} x^{-\varepsilon, \mathbf{0}}, \quad (2.101)$$

where $\alpha \in \ker_{\varphi_2} \setminus \{-\varepsilon\}$, and $u' \in \mathcal{A}_{\langle \hat{k}-1 \rangle} + \mathbb{F} x^{-\varepsilon, \mathbf{0}}$. In the following, we will use u' frequently in its specific form as in (2.83).

Subcase 2.1. There exists $q \in \overline{3, 2n}$ and $a_{\alpha, \mathbf{i}} \neq 0$ such that $i_q > 0$ and $\varphi_q(\alpha) = 0$.

If $\varphi_{q'} \neq 0$, choose $\tau' \in \text{Rad}_{\phi} \cap (\cap_{q' \neq p \in \overline{1, 2n}} \ker_{\varphi_p}) \setminus \ker_{\varphi_{q'}}$ such that

$$\varphi_{q'}(\alpha + \tau' + \sigma_q) \neq 0. \quad (2.102)$$

Then we have

$$\begin{aligned} [x^{\tau', \mathbf{0}}, u] &\equiv \left(\sum_{\substack{\mathbf{i} \in \mathcal{J} \\ i_2=0, |\mathbf{i}|=\hat{k}}} a_{\alpha, \mathbf{i}} \epsilon_{q'} i_q \varphi_{q'}(\tau') x^{\alpha + \tau' + \sigma_q + \varepsilon, \mathbf{i} - 1_{[q]}} \right. \\ &\quad \left. + \sum_{\substack{\alpha + \tau' + \sigma_q + \varepsilon \neq \beta \in \Gamma, \mathbf{l} \in \mathcal{J} \\ l_2=0, |\mathbf{l}|=\hat{k}-1}} c'_{\beta, \mathbf{l}} x^{\beta, \mathbf{l}} \right) \pmod{\mathcal{A}_{\langle \hat{k}-2 \rangle}} \end{aligned} \quad (2.103)$$

with $c'_{\beta, \mathbf{l}} \in \mathbb{F}$. Since $\varphi_{q'}(\alpha + \tau' + \sigma_q + \varepsilon) \neq 0$ by (2.102), we have $\alpha + \tau' + \sigma_q + \varepsilon \neq -\varepsilon$. Then $[x^{\tau', \mathbf{0}}, u] \in (\mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k}-1 \rangle}) \setminus \mathbb{F} x^{-\varepsilon, \mathbf{0}}$, which contradicts the minimality of $\hat{k}$. Assume that $\varphi_{q'} \equiv 0$, then $\mathcal{J}_{q'} = \mathbb{N}$. Picking $\tau \in \text{Rad}_{\phi} \cap (\cap_{p \in \overline{2, 2n}} \ker_{\varphi_p}) \setminus \ker_{\varphi_1}$ such that $\varphi_1(\alpha + \tau + \varepsilon) + 1 \neq 0$, we have

$$\begin{aligned} [x^{\tau, 1_{[q]}}, u] &\equiv \left(\sum_{\substack{\mathbf{i} \in \mathcal{J} \\ i_2=0, |\mathbf{i}|=\hat{k}}} a_{\alpha, \mathbf{i}} \epsilon_{q'} i_q x^{\alpha + \tau + \sigma_q + \varepsilon, \mathbf{i} - 1_{[q]}} \right. \\ &\quad \left. + \sum_{\substack{\alpha + \tau + \sigma_q + \varepsilon \neq \beta \in \Gamma, \mathbf{l} \in \mathcal{J} \\ l_2=0, |\mathbf{l}|=\hat{k}-1}} c'_{\beta, \mathbf{l}} x^{\beta, \mathbf{l}} \right) \pmod{\mathcal{A}_{\langle \hat{k}-2 \rangle}} \end{aligned} \quad (2.104)$$

with $c'_{\beta, \mathbf{l}} \in \mathbb{F}$. Since $\varphi_1(\alpha + \tau + \sigma_q + \varepsilon) + 1 \neq 0$, we have $\alpha + \tau + \sigma_q + \varepsilon \neq -\varepsilon$. Then $[x^{\tau, 1_{[q]}}, u] \in (\mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k}-1 \rangle}) \setminus \mathbb{F} x^{-\varepsilon, \mathbf{0}}$, which contradicts the minimality of $\hat{k}$.

Subcase 2.2. There exists $q \in \overline{3, 2n}$ and $a_{\alpha, \mathbf{i}} \neq 0$ such that $i_q > 0$ and $\varphi_q(\alpha) \neq 0$.

If $\varphi_{q'} \neq 0$, choose $\tau' \in \text{Rad}_{\phi} \cap (\cap_{q' \neq p \in \overline{1, 2n}} \ker_{\varphi_p}) \setminus \ker_{\varphi_{q'}}$ such that

$$\varphi_q(\alpha) \varphi_{q'}(\tau' - \sigma_q) + \varphi_{q'}(\alpha) \varphi_q(\sigma_q) \neq 0. \quad (2.105)$$

Then we have

$$\begin{aligned} (\mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle}) \setminus \mathbb{F} x^{-\varepsilon, \mathbf{0}} &\ni [x^{\tau' - \alpha - \sigma_q - \varepsilon, \mathbf{0}}, u] \equiv \sum_{\substack{\mathbf{i} \in \mathcal{J} \\ i_2=0, |\mathbf{i}|=\hat{k}}} a_{\alpha, \mathbf{i}} \epsilon_{q'} (\varphi_q(\alpha) \varphi_{q'}(\tau' - \sigma_q) \\ &\quad + \varphi_{q'}(\alpha) \varphi_q(\sigma_q)) x^{\tau', \mathbf{i}} \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}}. \end{aligned} \quad (2.106)$$

Since $\varphi_q(\tau') = 0$, we go back to *Subcase 2.1* with u replaced by $[x^{\tau' - \alpha - \sigma_q - \varepsilon, \mathbf{0}}, u]$. Assume that $\varphi_{q'} \equiv 0$, by (1.7) we have $\mathcal{J}_{q'} = \mathbb{N}$. Then we get

$$\begin{aligned}
& \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k}-1 \rangle} \ni [x^{-\alpha - \sigma_q, \mathbf{0}}, u] \\
& \equiv \sum_{\substack{\mathbf{i} \in \mathcal{J} \\ i_2=0, |\mathbf{i}|=\hat{k}}} \sum_{s=1}^{n-1} a_{\alpha, \mathbf{i}} (i_{2s+1} \varphi_{2s+2}(\alpha + \sigma_q) x^{\sigma_{2s+1} + \varepsilon - \sigma_q, \mathbf{i} - \mathbf{1}_{[2s+1]}} - i_{2s+2} \varphi_{2s+1}(\alpha + \sigma_q) \cdot \\
& \quad x^{\sigma_{2s+1} + \varepsilon - \sigma_q, \mathbf{i} - \mathbf{1}_{[2s+2]}}) + \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\beta, \mathbf{l}) \in \ker \varphi_2 \times \mathcal{J} \\ l_2=0, |\mathbf{l}|=\hat{k}-1}} c_{\beta, \mathbf{l}} \left(\sum_{s=1}^{n-1} (\varphi_{2s+1}(\beta) \varphi_{2s+2}(\alpha + \sigma_q) \right. \\
& \quad \left. - \varphi_{2s+1}(\alpha + \sigma_q) \varphi_{2s+2}(\beta)) x^{\beta + \sigma_{2s+1} + \varepsilon - \alpha - \sigma_q, \mathbf{l}} \right. \\
& \quad \left. - \phi(\alpha, \beta) x^{\beta + \varepsilon - \alpha - \sigma_q, \mathbf{l}} \right) \pmod{\mathcal{A}_{\langle \hat{k}-2 \rangle}} \tag{2.107}
\end{aligned}$$

and

$$\begin{aligned}
& \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k}-1 \rangle} \ni [x^{-\alpha - \sigma_q - \varepsilon, \mathbf{0}}, u] \\
& \equiv \sum_{\substack{\mathbf{i} \in \mathcal{J} \\ i_2=0, |\mathbf{i}|=\hat{k}}} \sum_{s=1}^{n-1} a_{\alpha, \mathbf{i}} (i_{2s+1} \varphi_{2s+2}(\alpha + \sigma_q) x^{\sigma_{2s+1} - \sigma_q, \mathbf{i} - \mathbf{1}_{[2s+1]}} - i_{2s+2} \varphi_{2s+1}(\alpha + \sigma_q) \cdot \\
& \quad \cdot x^{\sigma_{2s+1} - \sigma_q, \mathbf{i} - \mathbf{1}_{[2s+2]}}) + \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\beta, \mathbf{l}) \in \ker \varphi_2 \times \mathcal{J} \\ l_2=0, |\mathbf{l}|=\hat{k}-1}} c_{\beta, \mathbf{l}} \left(\sum_{s=1}^{n-1} (\varphi_{2s+1}(\beta) \varphi_{2s+2}(\alpha + \sigma_q) \right. \\
& \quad \left. - \varphi_{2s+1}(\alpha + \sigma_q) \varphi_{2s+2}(\beta)) x^{\beta + \sigma_{2s+1} - \alpha - \sigma_q, \mathbf{l}} \right. \\
& \quad \left. - \phi(\alpha, \beta) x^{\beta - \alpha - \sigma_q, \mathbf{l}} \right) \pmod{\mathcal{A}_{\langle \hat{k}-2 \rangle}}. \tag{2.108}
\end{aligned}$$

Since $\sigma_{2s+1} - \sigma_q \neq -\varepsilon$ for all $s \in \overline{1, n-1}$, avoiding contradiction to the minimality of $\hat{k}$, the coefficients corresponding to the terms $x^{\sigma_{2s+1} - \sigma_q, \mathbf{i} - \mathbf{1}_{[2s+1]}}$ and $x^{\sigma_{2s+1} - \sigma_q, \mathbf{i} - \mathbf{1}_{[2s+2]}}$ for all $s \in \overline{1, n-1}$ in (2.108) are zeros. For any $s \in \overline{1, n-1}$, we have

$$\beta + \sigma_{2s+1} + \varepsilon - \alpha - \sigma_q \neq -\varepsilon \text{ or } \beta + \sigma_{2s+1} - \alpha - \sigma_q \neq -\varepsilon. \tag{2.109}$$

The coefficient corresponding to the term $x^{\beta + \sigma_{2s+1} + \varepsilon - \alpha - \sigma_q, \mathbf{l}}$ in (2.107) and the coefficient corresponding to the term $x^{\beta + \sigma_{2s+1} - \alpha - \sigma_q, \mathbf{l}}$ in (2.108) are the same. To avoid contradiction to the minimality of $\hat{k}$, we get that these coefficients equal zero. Similarly, we have

$$\beta + \varepsilon - \alpha - \sigma_q \neq -\varepsilon \text{ or } \beta - \alpha - \sigma_q \neq -\varepsilon. \tag{2.110}$$

The coefficient corresponding to the term $x^{\beta+\varepsilon-\alpha-\sigma_q, \mathbf{1}}$ in (2.107) and the coefficient corresponding to the term $x^{\beta-\alpha-\sigma_q, \mathbf{1}}$ in (2.108) are the same. To avoid contradiction to the minimality of $\hat{k}$, we get that these coefficients equal zero too. Using these results, we have

$$\begin{aligned} \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle} &\ni [x^{-\alpha-\sigma_q-\varepsilon, \mathbf{1}_{[q]}}, u] \\ &\equiv \sum_{\substack{\mathbf{i} \in \mathcal{J} \\ i_2=0, |\mathbf{i}|=\hat{k}}} a_{\alpha, \mathbf{i}} \epsilon_{q'} \varphi_q(\alpha) x^{0, \mathbf{i}} \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}}. \end{aligned} \quad (2.111)$$

Thus we go back to *Subcase 2.1* with u replaced by $[x^{-\alpha-\sigma_q-\varepsilon, \mathbf{1}_{[q]}}, u]$.

Subcase 2.3. For any $a_{\alpha, \mathbf{i}} \neq 0$, $i_q = 0$ for all $q \in \overline{3, 2n}$.

Since $i_q = 0$ for all $q \in \overline{2, 2n}$, we obtain $i_1 = \hat{k}$. Then we have

$$u = a_{\alpha, \hat{k}_{[1]}} x^{\alpha, \hat{k}_{[1]}} + u' \in (\mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle}) \setminus \mathbb{F} x^{-\varepsilon, \mathbf{0}}. \quad (2.112)$$

If $\varphi_2 \neq 0$, pick $\tau \in \text{Rad}_\phi \cap (\cap_{2 \neq p \in \overline{1, 2n}} \ker \varphi_p) \setminus \ker \varphi_2$. Moreover, if $\varphi_1(\alpha) + 1 \neq 0$, we have

$$\begin{aligned} (\mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle}) \setminus \mathbb{F} x^{-\varepsilon, \mathbf{0}} &\ni [x^{-\tau-\varepsilon, \mathbf{0}}, [x^{\tau-\alpha, \mathbf{0}}, u]] \\ &\equiv -a_{\alpha, \hat{k}_{[1]}} \varphi_2^2(\tau) (\varphi_1(\alpha) + 1) x^{-\varepsilon, \hat{k}_{[1]}} \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}}. \end{aligned} \quad (2.113)$$

Since $\varphi_1(-\varepsilon) + 1 = 0$, replacing u by $[x^{-\tau-\varepsilon, \mathbf{0}}, [x^{\tau-\alpha, \mathbf{0}}, u]]$ and α by $-\varepsilon$, we can always assume that $\varphi_1(\alpha) + 1 = 0$. Choose $\tau' \in \text{Rad}_\phi \cap (\cap_{p \in \overline{2, 2n}} \ker \varphi_p) \setminus \ker \varphi_1$ such that $\varphi_1(\tau') + 1 \neq 0$. Then we have

$$\begin{aligned} [x^{-\tau+\tau', \mathbf{0}}, [x^{\tau, \mathbf{0}}, u]] &\equiv -a_{\alpha, \hat{k}_{[1]}} \hat{k} \varphi_2^2(\tau) (\varphi_1(\tau') + 1) x^{\alpha+\tau', (\hat{k}-1)_{[1]}} \\ &\quad + \sum_{\substack{\alpha+\tau' \neq \beta \in \Gamma, \mathbf{l} \in \mathcal{J} \\ l_2=0, |\mathbf{l}|=\hat{k}-1}} c'_{\beta, \mathbf{l}} x^{\beta, \mathbf{l}} \pmod{\mathcal{A}_{\langle \hat{k}-2 \rangle}} \end{aligned} \quad (2.114)$$

with $c'_{\beta, \mathbf{l}} \in \mathbb{F}$. Since $\varphi_1(\alpha + \tau') + 1 \neq 0$, we get $\alpha + \tau' \neq -\varepsilon$. Then we have $[x^{-\tau+\tau', \mathbf{0}}, [x^{\tau, \mathbf{0}}, u]] \in (\mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k}-1 \rangle}) \setminus \mathbb{F} x^{-\varepsilon, \mathbf{0}}$, which contradicts the minimality of $\hat{k}$.

Assume that $\varphi_2 \equiv 0$, by (1.7) we have $\mathcal{J}_2 = \mathbb{N}$. Then we get

$$\begin{aligned} \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k}-1 \rangle} &\ni [x^{-\alpha, \mathbf{0}}, u] \\ &\equiv \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\beta, \mathbf{l}) \in \ker \varphi_2 \times \mathcal{J} \\ l_2=0, |\mathbf{l}|=\hat{k}-1}} c_{\beta, \mathbf{l}} \left(\sum_{s=1}^{n-1} (\varphi_{2s+1}(\beta) \varphi_{2s+2}(\alpha) - \varphi_{2s+1}(\alpha) \varphi_{2s+2}(\beta)) \right) \cdot \\ &\quad x^{\beta+\sigma_{2s+1}+\varepsilon-\alpha, \mathbf{l}} - \phi(\alpha, \beta) x^{\beta+\varepsilon-\alpha, \mathbf{l}} \pmod{\mathcal{A}_{\langle \hat{k}-2 \rangle}} \end{aligned} \quad (2.115)$$

and

$$\begin{aligned}
& \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k}-1 \rangle} \ni [x^{-\alpha-\varepsilon, \mathbf{0}}, u] \\
& \equiv \sum_{\substack{(-\varepsilon, \mathbf{0}) \neq (\beta, \mathbf{l}) \in \ker \varphi_2 \times \mathcal{J} \\ l_2=0, |\mathbf{l}|=\hat{k}-1}} c_{\beta, \mathbf{l}} \left(\sum_{s=1}^{n-1} (\varphi_{2s+1}(\beta) \varphi_{2s+2}(\alpha) - \varphi_{2s+1}(\alpha) \varphi_{2s+2}(\beta)) \cdot \right. \\
& \quad \left. x^{\beta+\sigma_{2s+1}-\alpha, \mathbf{l}} - \phi(\alpha, \beta) x^{\beta-\alpha, \mathbf{l}} \right) \pmod{\mathcal{A}_{\langle \hat{k}-2 \rangle}}. \tag{2.116}
\end{aligned}$$

For any $s \in \overline{1, n-1}$, we have

$$\beta + \sigma_{2s+1} + \varepsilon - \alpha \neq -\varepsilon \text{ or } \beta + \sigma_{2s+1} - \alpha \neq -\varepsilon. \tag{2.117}$$

The coefficient corresponding to the term $x^{\beta+\sigma_{2s+1}+\varepsilon-\alpha, \mathbf{l}}$ in (2.115) and the coefficient corresponding to the term $x^{\beta+\sigma_{2s+1}-\alpha, \mathbf{l}}$ in (2.116) are the same. To avoid contradiction to the minimality of $\hat{k}$, we get that these coefficients equal zero. Similarly, we have

$$\beta + \varepsilon - \alpha \neq -\varepsilon \text{ or } \beta - \alpha \neq -\varepsilon. \tag{2.118}$$

The coefficient corresponding to the term $x^{\beta+\varepsilon-\alpha, \mathbf{l}}$ in (2.115) and the coefficient corresponding to the term $x^{\beta-\alpha, \mathbf{l}}$ in (2.116) are the same. To avoid contradiction to the minimality of $\hat{k}$, we get that these coefficients equal zero too. Using these results, we have

$$\begin{aligned}
& \mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k} \rangle} \ni [x^{-\alpha-\varepsilon, \mathbf{1}_{[2]}}, u] \\
& \equiv -a_{\alpha, \hat{k}_{[1]}} (\varphi_1(\alpha) + 1) x^{-\varepsilon, \hat{k}_{[1]}} \pmod{\mathcal{A}_{\langle \hat{k}-1 \rangle}}. \tag{2.119}
\end{aligned}$$

If $\varphi_1(\alpha) + 1 \neq 0$, by replacing u by $[x^{-\alpha-\varepsilon, \mathbf{1}_{[2]}}, u]$ and α by $-\varepsilon$, we can always assume that $\varphi_1(\alpha) + 1 = 0$. Picking $\tau' \in \text{Rad}_\phi \cap (\cap_{p \in \overline{2, 2n}} \ker \varphi_p) \setminus \ker \varphi_1$, we have

$$[x^{\tau', \mathbf{1}_{[2]}}, u] \equiv -\hat{k} a_{\alpha, \hat{k}_{[1]}} x^{\alpha+\tau', (\hat{k}-1)_{[1]}} + \sum_{\substack{\alpha+\tau' \neq \beta \in \Gamma, \mathbf{l} \in \mathcal{J} \\ l_2=0, |\mathbf{l}|=\hat{k}-1}} c'_{\beta, \mathbf{l}} x^{\beta, \mathbf{l}} \pmod{\mathcal{A}_{\langle \hat{k}-2 \rangle}} \tag{2.120}$$

with $c'_{\beta, \mathbf{l}} \in \mathbb{F}$. Since $\varphi_1(\alpha + \tau') + 1 = \varphi_1(\tau') \neq 0$, we have $\alpha + \tau' \neq -\varepsilon$. Then $[x^{\tau', \mathbf{1}_{[2]}}, u] \in (\mathcal{I}_{[0]} \cap \mathcal{A}_{\langle \hat{k}-1 \rangle}) \setminus \mathbb{F} x^{-\varepsilon, \mathbf{0}}$, which contradicts the minimality of $\hat{k}$.

So we always have $\hat{k} = 0$ and $1 \in \mathcal{I}$.

Step 3. The conclusion of Step 2 implies $\mathcal{I} = \mathcal{A}$ if $\mathcal{J} \neq \{0\}$.

Since

$$[1, x^{\beta, \mathbf{j}}] = \varphi_2(\beta) x^{\beta, \mathbf{j}} + j_2 x^{\beta, \mathbf{j}-1_{[2]}} \quad \text{for } (\beta, \mathbf{j}) \in \Gamma \times \mathcal{J}. \tag{2.121}$$

If $\mathcal{J}_2 = \mathbb{N}$, we can prove $\mathcal{I} = \mathcal{A}$ by induction on j_2 . Assume that $\mathcal{J}_2 = \{0\}$, then $\varphi_2 \not\equiv 0$ by (1.7). Thus (2.121) implies

$$x^{\beta, \mathbf{j}} \in \mathcal{I} \quad \text{for } (\beta, \mathbf{j}) \in \Gamma \times \mathcal{J}, \varphi_2(\beta) \neq 0. \quad (2.122)$$

Picking $\tau \in \text{Rad}_\phi \cap (\cap_{2 \neq p \in \overline{1, 2n}} \ker_{\varphi_p}) \setminus \ker_{\varphi_2}$, we get $x^{\tau, \mathbf{0}} \in \mathcal{I}$. Since we have

$$\mathcal{I} \ni [x^{\tau, \mathbf{0}}, x^{-\tau + \beta, \mathbf{j}}] = -\varphi_2(\tau)(\varphi_1(\beta) + 2)x^{\beta, \mathbf{j}} - j_1\varphi_2(\tau)x^{\beta, \mathbf{j}-1[1]} \quad (2.123)$$

for any $(\beta, \mathbf{j}) \in \ker_{\varphi_2} \times \mathcal{J}$. If $\mathcal{J}_1 = \mathbb{N}$, we can prove $\mathcal{I} = \mathcal{A}$ by induction on j_1 . Assume that $\mathcal{J}_1 = \{0\}$. Then (2.123) shows

$$x^{\beta, \mathbf{j}} \in \mathcal{I} \quad \text{for } (\beta, \mathbf{j}) \in \ker_{\varphi_2} \times \mathcal{J}, \varphi_1(\beta) + 2 \neq 0. \quad (2.124)$$

If there exists $q \in \overline{3, 2n}$ such that $\mathcal{J}_q \neq \{0\}$. Moreover, if $\varphi_{q'} \not\equiv 0$, picking $\tau \in \text{Rad}_\phi \cap (\cap_{q' \neq p \in \overline{1, 2n}} \ker_{\varphi_p}) \setminus \ker_{\varphi_{q'}}$, we have $x^{\tau, \mathbf{0}} \in \mathcal{I}$ due to $\varphi_1(\tau) + 2 \neq 0$. Since we get

$$\mathcal{I} \ni [x^{\tau, \mathbf{0}}, x^{\beta - \tau - \sigma_q - \varepsilon, \mathbf{j}}] = \epsilon_{q'}(\varphi_{q'}(\tau)\varphi_q(\beta - \sigma_q)x^{\beta, \mathbf{j}} + j_q\varphi_{q'}(\tau)x^{\beta, \mathbf{j}-1[q]}) \quad (2.125)$$

for $\beta \in \ker_{\varphi_2}$. Then by (2.125) and induction on j_q , we have $\mathcal{I} = \mathcal{A}$. Assume that $\varphi_{q'} \equiv 0$, then $\mathcal{J}_{q'} = \mathbb{N}$. Since $x^{-\sigma_q - \varepsilon, 1[q]} \in \mathcal{I}$ due to $\varphi_1(-\sigma_q - \varepsilon) + 2 \neq 0$, we have

$$\mathcal{I} \ni [x^{-\sigma_q - \varepsilon, 1[q]}, x^{\beta, \mathbf{j}}] = \epsilon_{q'}((j_{q'}\varphi_q(\sigma_q) + \varphi_q(\beta))x^{\beta, \mathbf{j}} + j_qx^{\beta, \mathbf{j}-1[q]}) \quad (2.126)$$

for $\beta \in \ker_{\varphi_2}$. Then by (2.126) and induction on j_q , we have $\mathcal{I} = \mathcal{A}$.

Therefore, we have proved that $\mathcal{I} = \mathcal{A}$ if $\mathcal{J} \neq \{0\}$. So the first statement of Theorem 1 holds.

Step 4. The second statement of Theorem 1 holds.

If $\mathcal{J} = \{0\}$, we have $\varphi_p \not\equiv 0$ for any $p \in \overline{1, 2n}$. Fixing any $q \in \overline{3, 2n}$, we pick $\tau \in \text{Rad}_\phi \cap (\cap_{q' \neq p \in \overline{1, 2n}} \ker_{\varphi_p}) \setminus \ker_{\varphi_{q'}}$. Since $\varphi_1(\tau) + 2 \neq 0$, we have $x^{\tau, \mathbf{0}} \in \mathcal{I}$ by (2.124). Then we get

$$\mathcal{I} \ni [x^{\tau, \mathbf{0}}, x^{\beta - \tau - \sigma_q - \varepsilon, \mathbf{0}}] = \epsilon_{q'}\varphi_{q'}(\tau)\varphi_q(\beta - \sigma_q)x^{\beta, \mathbf{0}} \quad (2.127)$$

for $\beta \in \ker_{\varphi_2}$. Thus we obtain

$$x^{\beta, \mathbf{0}} \in \mathcal{I}, \quad \text{for } \beta \in \ker_{\varphi_2} \text{ such that } \varphi_q(\beta - \sigma_q) \neq 0. \quad (2.128)$$

For $\beta \in \ker_{\varphi_2}$ such that $\varphi_1(\beta) + 2 = 0$ and $\varphi_q(\beta - \sigma_q) = 0$ for all $q \in \overline{3, 2n}$, if $\beta \notin \text{Rad}_\phi$, there exists $\alpha \in \Gamma$ such that $\phi(\alpha, \beta) \neq 0$. Fix any such α . If

$\varphi_1(\alpha) + 2 = 0$, replace α by $\alpha + \varepsilon$. Then we can always assume that $\varphi_1(\alpha) + 2 \neq 0$. By (2.122) and (2.124), we have $x^{\alpha, \mathbf{0}} \in \mathcal{I}$. Then we get

$$\begin{aligned} \mathcal{I} \ni [x^{\alpha, \mathbf{0}}, x^{\beta - \alpha - \varepsilon, \mathbf{0}}] &= \sum_{s=1}^{n-1} (\varphi_{2s+1}(\alpha) \varphi_{2s+2}(\beta) - \varphi_{2s+2}(\alpha) \varphi_{2s+1}(\beta)) x^{\beta + \sigma_{2s+1}, \mathbf{0}} \\ &\quad + \phi(\alpha, \beta) x^{\beta, \mathbf{0}} - (\varphi_1(\beta) + 1) \varphi_2(\alpha) x^{\beta - \varepsilon, \mathbf{0}}. \end{aligned} \quad (2.129)$$

Since $\varphi_1(\beta - \varepsilon) + 2 \neq 0$, we have $x^{\beta - \varepsilon, \mathbf{0}} \in \mathcal{I}$. Since

$$\varphi_{2s+1}(\beta + \sigma_{2s+1} - \sigma_{2s+1}) = \varphi_{2s+1}(\sigma_{2s+1}) \neq 0 \quad (2.130)$$

or

$$\varphi_{2s+2}(\beta + \sigma_{2s+1} - \sigma_{2s+2}) = \varphi_{2s+2}(\sigma_{2s+1}) \neq 0, \quad (2.131)$$

we have $x^{\beta + \sigma_{2s+1}, \mathbf{0}} \in \mathcal{I}$ for all $s \in \overline{1, n-1}$. Then by (2.129) we get

$$x^{\beta, \mathbf{0}} \in \mathcal{I} \text{ for } \beta \in \ker_{\varphi_2} \setminus \text{Rad}_{\phi}, \varphi_1(\beta) + 2 = 0, \varphi_q(\beta - \sigma_q) = 0, q \in \overline{3, 2n}. \quad (2.132)$$

Thus we have obtained that

$$x^{\beta, \mathbf{0}} \in \mathcal{I} \text{ for } \beta \in \Gamma \setminus \{\sigma\} \quad (2.133)$$

by (2.122), (2.124), (2.128) and (2.132). Set

$$\hat{\mathcal{H}} = \text{span}\{x^{\alpha, \mathbf{0}} \mid \alpha \in \Gamma \setminus \{\sigma\}\}. \quad (2.134)$$

Since $\mathcal{J} = \{0\}$, by (2.1), (2.121), (2.123), (2.127) and (2.129), we have

$$\hat{\mathcal{H}} = [\mathcal{A}, \mathcal{A}]. \quad (2.135)$$

Replacing $\mathcal{A}$ by $\hat{\mathcal{H}}$ and Γ by $\Gamma \setminus \{\sigma\}$ in the above proof, we can also obtain (2.133), which implies the simplicity of

$$\mathcal{H}^{(1)} = [\mathcal{H}, \mathcal{H}] = \hat{\mathcal{H}} / \mathbb{F}x^{-\varepsilon, \mathbf{0}}.$$

This completes the proof of Theorem 1. $\square$

3 Proof of Theorem 2

In this section, we will determine the irreducibility of the module $\mathcal{A}_{\xi, f}$.

If there exists $\mu \in \Gamma$ such that (1.16) and (1.17) hold, by (1.15) and (2.1) we have the module isomorphism from $\mathcal{A}_{\xi,f}$ to $\mathcal{A}_{\xi_0,0}$ in the following:

$$\psi : \mathcal{A}_{\xi,f} \longrightarrow \mathcal{A}_{\xi_0,0}, \quad (3.1)$$

$$x^{\beta,\mathbf{j}} \longmapsto x^{\beta+\mu-\varepsilon,\mathbf{j}}. \quad (3.2)$$

As a $\mathcal{H}$ -module, $\mathcal{A}_{\xi_0,0}$ has a one dimensional trivial submodule $\mathbb{F}x^{-\varepsilon,\mathbf{0}}$. And $\mathcal{A}_{\xi_0,0}/\mathbb{F}x^{-\varepsilon,\mathbf{0}}$ is irreducible when $\mathcal{J} \neq \{\mathbf{0}\}$ and indecomposable when $\mathcal{J} = \{\mathbf{0}\}$.

In the following, we assume that such μ does not exist.

Let $\mathcal{N}$ be any nonzero submodule of $\mathcal{A}_{\xi,f}$. To prove the second statement of Theorem 2 is equivalent to proving that $\mathcal{N} = \mathcal{A}_{\xi,f}$.

For $k \in \mathbb{N}$, let

$$\mathcal{A}_{\xi,f}^{(k)} = \text{span}\{x^{\alpha,\mathbf{i}} \mid (\alpha,\mathbf{i}) \in \Gamma \times \mathcal{J}, |\mathbf{i}| \leq k\}. \quad (3.3)$$

Then we have

$$\mathcal{A}_{\xi,f} = \bigcup_{k=0}^{\infty} \mathcal{A}_{\xi,f}^{(k)}. \quad (3.4)$$

Set

$$\hat{k}' = \min\{k \in \mathbb{N} \mid \mathcal{N} \cap \mathcal{A}_{\xi,f}^{(k)} \neq \{0\}\}. \quad (3.5)$$

Then for any $u \in (\mathcal{N} \cap \mathcal{A}_{\xi,f}^{(\hat{k}')}) \setminus \{0\}$, we write it as

$$u = \sum_{\substack{(\alpha,\mathbf{i}) \in \Gamma \times \mathcal{J}, \\ |\mathbf{i}| = \hat{k}'}} b_{\alpha,\mathbf{i}} x^{\alpha,\mathbf{i}} + u', \quad (3.6)$$

where $b_{\alpha,\mathbf{i}} \in \mathbb{F}$ and $u' \in \mathcal{A}_{\xi,f}^{(\hat{k}'-1)}$. Moreover, we define

$$\kappa(u) = |\{\alpha \in \Gamma \mid b_{\alpha,\mathbf{i}} \neq 0 \text{ for some } \mathbf{i} \in \mathcal{J}, |\mathbf{i}| = \hat{k}'\}|. \quad (3.7)$$

By (3.5), $\kappa(u) > 0$. Furthermore, we set

$$\kappa = \min\{\kappa(w) \mid w \in (\mathcal{N} \cap \mathcal{A}_{\xi,f}^{(\hat{k}')}) \setminus \{0\}\}. \quad (3.8)$$

Choose $u \in (\mathcal{N} \cap \mathcal{A}_{\xi,f}^{(\hat{k}')}) \setminus \{0\}$ such that $\kappa(u) = \kappa$. Write u as in (3.6).

Quite similarly to the discussion in Step 1 and the second case of Step 2 of the proof of Theorem 1, we can obtain some $x^{\alpha,\mathbf{0}} \in \mathcal{N}$ with $\alpha \in \Gamma$. Here we omit

the details and let α be fixed. Then we will prove $\mathcal{N} = \mathcal{A}_{\xi, f}$ by induction on $|\mathbf{i}|$. Frequently, we will use the notation of Lie algebra homomorphism

$$\rho : \mathcal{H} \longrightarrow \mathrm{GL}(\mathcal{A}_{\xi, f}) \quad (3.9)$$

to denote the module action in the rest of this paper.

Firstly we will prove $x^{\gamma, \mathbf{i}} \in \mathcal{N}$ for all $\gamma \in \Gamma$ when $|\mathbf{i}| = 0$. Namely, we will prove $x^{\gamma, \mathbf{0}} \in \mathcal{N}$ for all $\gamma \in \Gamma$. We give the proof in four cases.

Case 1. $\gamma \in \Gamma$ such that

$$(\varphi_1(\gamma) + 1 + \xi_1)(\varphi_2(\alpha) + \xi_2) - (\varphi_2(\gamma) + \xi_2)(\varphi_1(\alpha) + \xi_1) \neq 0. \quad (3.10)$$

By (1.15), we have

$$\begin{aligned} \mathcal{N} &\ni \bar{x}^{\gamma-\alpha, \mathbf{0}}.x^{\alpha, \mathbf{0}} \\ &= ((\varphi_1(\gamma) + 1 + \xi_1)(\varphi_2(\alpha) + \xi_2) - (\varphi_2(\gamma) + \xi_2)(\varphi_1(\alpha) + \xi_1))x^{\gamma, \mathbf{0}} \\ &\quad + \sum_{s=1}^{n-1} ((\varphi_{2s+1}(\gamma) + \xi_{2s+1})(\varphi_{2s+2}(\alpha) + \xi_{2s+2}) - (\varphi_{2s+2}(\gamma) + \xi_{2s+2}) \\ &\quad \cdot (\varphi_{2s+1}(\alpha) + \xi_{2s+1}))x^{\gamma+\sigma_{2s+1}+\varepsilon, \mathbf{0}} + (\phi(\gamma, \alpha) + f(\gamma - \alpha))x^{\gamma+\varepsilon, \mathbf{0}}. \end{aligned} \quad (3.11)$$

If $\varphi_2 \equiv 0$, we have $\mathcal{J}_2 = \mathbb{N}$. Moreover, if $\xi_2 = 0$, there does not exist any $\gamma \in \Gamma$ such that (3.10) holds. We assume that $\xi_2 \neq 0$. Then we have

$$\bar{x}^{\varepsilon, \mathbf{0}}.(\bar{x}^{-\varepsilon, 1[2]}.x^{\beta, \mathbf{0}}) = -2\xi_2(\varphi_1(\beta) + \xi_1)x^{\beta, \mathbf{0}}. \quad (3.12)$$

Repeatedly applying $\rho(\bar{x}^{\varepsilon, \mathbf{0}})\rho(\bar{x}^{-\varepsilon, 1[2]})$ on $\mathcal{N}$, we obtain $x^{\gamma, \mathbf{0}} \in \mathcal{N}$ by Lemma 3, (3.10), (3.11) and (3.12). If $\varphi_2 \not\equiv 0$, picking $\tau \in \mathrm{Rad}_\phi \cap (\cap_{2 \neq p \in 1, 2n} \overline{\ker_{\varphi_p}}) \setminus \ker_{\varphi_2}$, we have

$$\bar{x}^{\varepsilon-\tau, \mathbf{0}}.(\bar{x}^{\tau-\varepsilon, \mathbf{0}}.x^{\beta, \mathbf{0}}) = S_1(\beta)x^{\beta, \mathbf{0}}, \quad (3.13)$$

where $S_1(\beta) = -\varphi_2(\tau)(\varphi_1(\beta) + \xi_1)(2(\varphi_2(\beta) + \xi_2) + \varphi_2(\tau)(\varphi_1(\beta) + \xi_1 + 1))$. Since we have

$$S_1(\gamma + \varepsilon) - S_1(\gamma) = -2\varphi_2(\tau)(\varphi_2(\tau)(\varphi_1(\gamma) + 1 + \xi_1) + \varphi_2(\gamma) + \xi_2), \quad (3.14)$$

and by (3.10) we have $\varphi_1(\gamma) + 1 + \xi_1 \neq 0$ or $\varphi_2(\gamma) + \xi_2 \neq 0$, we can restrict the selection of τ so that $S_1(\gamma + \varepsilon) - S_1(\gamma) \neq 0$. Then repeatedly applying $\rho(\bar{x}^{\varepsilon-\tau, \mathbf{0}})\rho(\bar{x}^{\tau-\varepsilon, \mathbf{0}})$ on $\mathcal{N}$, we obtain that $x^{\gamma, \mathbf{0}} \in \mathcal{N}$ by Lemma 3, (3.10), (3.11) and (3.13).

Case 2. $\gamma \in \Gamma$ such that

$$\begin{aligned} & (\varphi_{2t+1}(\gamma - \sigma_{2t+1}) + \xi_{2t+1})(\varphi_{2t+2}(\alpha) + \xi_{2t+2}) \\ & - (\varphi_{2t+2}(\gamma - \sigma_{2t+1}) + \xi_{2t+2})(\varphi_{2t+1}(\alpha) + \xi_{2t+1}) \neq 0 \end{aligned} \quad (3.15)$$

for some $t \in \overline{1, n-1}$.

By (1.15), we have

$$\begin{aligned} \mathcal{N} & \ni \bar{x}^{\gamma - \sigma_{2t+1} - \varepsilon - \alpha, \mathbf{0}} . x^{\alpha, \mathbf{0}} \\ & = ((\varphi_1(\gamma) + \xi_1)(\varphi_2(\alpha) + \xi_2) - (\varphi_2(\gamma) + \xi_2)(\varphi_1(\alpha) + \xi_1))x^{\gamma - \sigma_{2t+1} - \varepsilon, \mathbf{0}} \\ & \quad + \sum_{\substack{s=1 \\ s \neq t}}^{n-1} ((\varphi_{2s+1}(\gamma) + \xi_{2s+1})(\varphi_{2s+2}(\alpha) + \xi_{2s+2}) - (\varphi_{2s+2}(\gamma) + \xi_{2s+2}) \\ & \quad \cdot (\varphi_{2s+1}(\alpha) + \xi_{2s+1}))x^{\gamma + \sigma_{2s+1} - \sigma_{2t+1}, \mathbf{0}} + (\phi(\gamma, \alpha) + f(\gamma - \alpha))x^{\gamma - \sigma_{2t+1}, \mathbf{0}} \\ & \quad + ((\varphi_{2t+1}(\gamma - \sigma_{2t+1}) + \xi_{2t+1})(\varphi_{2t+2}(\alpha) + \xi_{2t+2}) - (\varphi_{2t+1}(\alpha) + \xi_{2t+1}) \\ & \quad \cdot (\varphi_{2t+2}(\gamma - \sigma_{2t+1}) + \xi_{2t+2}))x^{\gamma, \mathbf{0}}. \end{aligned} \quad (3.16)$$

If $\sigma_{2t+1} = 0$, we have $\varphi_{2t+1} \equiv \varphi_{2t+2} \equiv 0$ by (1.10). Then there does not exist any $\gamma \in \Gamma$ such that (3.15) holds. Assume that $\sigma_{2t+1} \neq 0$, we have $\varphi_{2t+1}(\sigma_{2t+1}) \neq 0$ or $\varphi_{2t+2}(\sigma_{2t+1}) \neq 0$ by (1.10). Choose $q \in \{2t+1, 2t+2\}$ such that $\varphi_q(\sigma_{2t+1}) \neq 0$. Then $\varphi_q \neq 0$, and we have that $\sigma_q = \sigma_{q'} = \sigma_{2t+1}$.

If $\varphi_{q'} \equiv 0$, we have $\mathcal{J}_{q'} = \mathbb{N}$. Moreover, if $\xi_{q'} = 0$, there does not exist any $\gamma \in \Gamma$ such that (3.15) holds. Suppose that $\xi_{q'} \neq 0$. Then we have

$$\bar{x}^{-2\sigma_q - \varepsilon, \mathbf{0}} . (\bar{x}^{-\varepsilon, 1_{[q']}} . x^{\beta, \mathbf{0}}) = 2\xi_{q'}\varphi_q(\sigma_q)(\varphi_q(\beta) + \xi_q)x^{\beta, \mathbf{0}}. \quad (3.17)$$

Repeatedly applying $\rho(\bar{x}^{-2\sigma_q - \varepsilon, \mathbf{0}})\rho(\bar{x}^{-\varepsilon, 1_{[q']}})$ on $\mathcal{N}$, we obtain that $x^{\gamma, \mathbf{0}} \in \mathcal{N}$ by Lemma 3, (3.15), (3.16) and (3.17).

If $\varphi_{q'} \neq 0$, picking $\tau' \in \text{Rad}_\phi \cap (\cap_{q' \neq p \in \overline{1, 2n}} \ker_{\varphi_p}) \setminus \ker_{\varphi_{q'}}$, we have

$$\bar{x}^{-\tau' - 2\sigma_q - \varepsilon, \mathbf{0}} . (\bar{x}^{\tau' - \varepsilon, \mathbf{0}} . x^{\beta, \mathbf{0}}) = S_2(\beta)x^{\beta, \mathbf{0}}, \quad (3.18)$$

where

$$\begin{aligned} S_2(\beta) & = \varphi_{q'}(\tau')(\varphi_q(\beta) + \xi_q)(2\varphi_q(\sigma_q)(\varphi_{q'}(\beta + \tau' + \sigma_q) + \xi_{q'}) \\ & \quad - \varphi_{q'}(\tau' + 2\sigma_q)(\varphi_q(\beta + \sigma_q) + \xi_q)). \end{aligned} \quad (3.19)$$

Since

$$\begin{aligned} S_2(\gamma) - S_2(\gamma - \sigma_q) & = 2\varphi_{q'}(\tau')\varphi_q(\sigma_q)(\varphi_q(\sigma_q)(\varphi_{q'}(\gamma - \sigma_q) + \xi_{q'}) \\ & \quad - \varphi_{q'}(\tau' + \sigma_q)(\varphi_q(\gamma - \sigma_q) + \xi_q)), \end{aligned} \quad (3.20)$$

and by (3.15) we have $\varphi_q(\gamma - \sigma_q) + \xi_q \neq 0$ or $\varphi_{q'}(\gamma - \sigma_q) + \xi_{q'} \neq 0$, we can restrict the selection of τ' so that $S_2(\gamma) - S_2(\gamma - \sigma_q) \neq 0$. Then repeatedly applying $\rho(\bar{x}^{-\tau' - 2\sigma_q - \varepsilon, \mathbf{0}})\rho(\bar{x}^{\tau' - \varepsilon, \mathbf{0}})$ on $\mathcal{N}$, we obtain that $x^{\gamma, \mathbf{0}} \in \mathcal{N}$ by Lemma 3, (3.15), (3.16) and (3.18).

Case 3. $\gamma \in \Gamma$ such that

$$\phi(\gamma, \alpha) + f(\gamma - \alpha) \neq 0. \quad (3.21)$$

By (1.15), we have

$$\begin{aligned} \mathcal{N} &\ni \bar{x}^{\gamma - \varepsilon - \alpha, \mathbf{0}}.x^{\alpha, \mathbf{0}} \\ &= ((\varphi_1(\gamma) + \xi_1)(\varphi_2(\alpha) + \xi_2) - (\varphi_2(\gamma) + \xi_2)(\varphi_1(\alpha) + \xi_1))x^{\gamma - \varepsilon, \mathbf{0}} \\ &\quad + \sum_{s=1}^{n-1} ((\varphi_{2s+1}(\gamma) + \xi_{2s+1})(\varphi_{2s+2}(\alpha) + \xi_{2s+2}) - (\varphi_{2s+2}(\gamma) + \xi_{2s+2}) \\ &\quad \cdot (\varphi_{2s+1}(\alpha) + \xi_{2s+1}))x^{\gamma + \sigma_{2s+1}, \mathbf{0}} + (\phi(\gamma, \alpha) + f(\gamma - \alpha))x^{\gamma, \mathbf{0}}. \end{aligned} \quad (3.22)$$

Similarly to the discussion from (3.11) to (3.18), we can obtain that $x^{\gamma, \mathbf{0}} \in \mathcal{N}$.

Case 4. $\gamma \in \Gamma$ such that

$$(\varphi_1(\gamma) + 1 + \xi_1)(\varphi_2(\alpha) + \xi_2) - (\varphi_2(\gamma) + \xi_2)(\varphi_1(\alpha) + \xi_1) = 0, \quad (3.23)$$

$$\begin{aligned} &(\varphi_{2t+1}(\gamma - \sigma_{2t+1}) + \xi_{2t+1})(\varphi_{2t+2}(\alpha) + \xi_{2t+2}) \\ &- (\varphi_{2t+2}(\gamma - \sigma_{2t+1}) + \xi_{2t+2})(\varphi_{2t+1}(\alpha) + \xi_{2t+1}) = 0 \end{aligned} \quad (3.24)$$

for all $t \in \overline{1, n-1}$, and

$$\phi(\gamma, \alpha) + f(\gamma - \alpha) = 0. \quad (3.25)$$

Subcase 4.1. $\varphi_2(\gamma) + \xi_2 \neq 0$.

Choose $\tau \in \text{Rad}_\phi \cap (\cap_{p \in \overline{2, 2n}} \ker \varphi_p) \setminus \ker \varphi_1$ such that $\varphi_1(\tau) + 1 \neq 0$.

If $\varphi_2(\alpha) + \xi_2 \neq 0$, we have

$$\mathcal{N} \ni \bar{x}^{\tau, \mathbf{0}}.x^{\alpha, \mathbf{0}} = (\varphi_1(\tau) + 1)(\varphi_2(\alpha) + \xi_2)x^{\alpha + \tau, \mathbf{0}}. \quad (3.26)$$

Then we go back to *Case 1* with $x^{\alpha, \mathbf{0}}$ replaced by $x^{\alpha + \tau, \mathbf{0}}$.

If $\varphi_2(\alpha) + \xi_2 = 0$, by (3.23) we have $\varphi_1(\alpha) + \xi_1 = 0$.

Moreover, if there exists $q \in \overline{3, 2n}$ such that $\varphi_q(\alpha) + \xi_q \neq 0$ and $\varphi_{q'} \not\equiv 0$, picking $\tau' \in \text{Rad}_\phi \cap (\cap_{q' \neq p \in \overline{1, 2n}} \ker \varphi_p) \setminus \ker \varphi_{q'}$, we have

$$\mathcal{N} \ni \bar{x}^{\tau + \tau', \mathbf{0}}.x^{\alpha, \mathbf{0}} = \epsilon_{q'} \varphi_{q'}(\tau')(\varphi_q(\alpha) + \xi_q)x^{\alpha + \tau + \tau' + \sigma_q + \varepsilon, \mathbf{0}}. \quad (3.27)$$

Then we go back to *Case 1* with $x^{\alpha, \mathbf{0}}$ replaced by $x^{\alpha+\tau+\tau'+\sigma_q+\varepsilon, \mathbf{0}}$.

If there exists $q \in \overline{3, 2n}$ such that $\varphi_q(\alpha) + \xi_q \neq 0$ and $\varphi_{q'} \equiv 0$, we have

$$\mathcal{N} \ni \bar{x}^{\tau, 1_{[q']}}.x^{\alpha, \mathbf{0}} = \epsilon_{q'}(\varphi_q(\alpha) + \xi_q)x^{\alpha+\tau+\sigma_q+\varepsilon, \mathbf{0}}. \quad (3.28)$$

We go back to *Case 1* with $x^{\alpha, \mathbf{0}}$ replaced by $x^{\alpha+\tau+\sigma_q+\varepsilon, \mathbf{0}}$.

If $\varphi_q(\alpha) + \xi_q = 0$ for all $q \in \overline{3, 2n}$, then there exists $\theta \in \Gamma$ such that

$$\phi(\theta, \alpha) + f(\theta) \neq 0. \quad (3.29)$$

We rechoose $\tau \in \text{Rad}_\phi \cap (\cap_{p \in \overline{2, 2n}} \ker_{\varphi_p}) \setminus \ker_{\varphi_1}$ such that $\varphi_1(\alpha + \theta + \tau + \varepsilon) + \xi_1 \neq 0$. Then we have

$$\mathcal{N} \ni \bar{x}^{\theta+\tau, \mathbf{0}}.x^{\alpha, \mathbf{0}} = (\phi(\theta, \alpha) + f(\theta))x^{\alpha+\theta+\tau+\varepsilon, \mathbf{0}}. \quad (3.30)$$

We go back to *Case 1* or the first condition of *Subcase 4.1* with $x^{\alpha, \mathbf{0}}$ replaced by $x^{\alpha+\theta+\tau+\varepsilon, \mathbf{0}}$. So we obtain that $x^{\gamma, \mathbf{0}} \in \mathcal{N}$.

Subcase 4.2. $\varphi_2(\gamma) + \xi_2 = 0$ and $\varphi_1(\gamma) + 1 + \xi_1 \neq 0$.

By (3.23) we have that $\varphi_2(\alpha) + \xi_2 = 0$. If $\varphi_2 \not\equiv 0$, we can prove $x^{\gamma, \mathbf{0}} \in \mathcal{N}$ similarly to *Subcase 4.1*. Here we omit the details. We assume that $\varphi_2 \equiv 0$. Then we get $\mathcal{J}_2 = \mathbb{N}$ and $\xi_2 = 0$.

If $\varphi_1(\alpha) + \xi_1 \neq 0$, we have

$$\begin{aligned} & \mathcal{N} \ni \bar{x}^{\gamma-\alpha, 1_{[2]}}.x^{\alpha, \mathbf{0}} \\ &= \sum_{s=1}^{n-1} ((\varphi_{2s+1}(\gamma) + \xi_{2s+1})(\varphi_{2s+2}(\alpha) + \xi_{2s+2}) - (\varphi_{2s+2}(\gamma) + \xi_{2s+2}) \\ & \quad \cdot (\varphi_{2s+1}(\alpha) + \xi_{2s+1}))x^{\gamma+\sigma_{2s+1}+\varepsilon, 1_{[2]}} - (\varphi_1(\alpha) + \xi_1)x^{\gamma, \mathbf{0}}. \end{aligned} \quad (3.31)$$

Since we have

$$\begin{aligned} & \bar{x}^{\varepsilon, 1_{[2]}}.(\bar{x}^{-\varepsilon, 1_{[2]}}.x^{\gamma+\sigma_{2s+1}+\varepsilon, 1_{[2]}}) \\ &= (\varphi_1(\gamma) + \xi_1 + 1)(\varphi_1(\gamma) + \xi_1 - 2)x^{\gamma+\sigma_{2s+1}+\varepsilon, 1_{[2]}} \end{aligned} \quad (3.32)$$

for $s \in \overline{1, n-1}$, and

$$\bar{x}^{\varepsilon, 1_{[2]}}.(\bar{x}^{-\varepsilon, 1_{[2]}}.x^{\gamma, \mathbf{0}}) = (\varphi_1(\gamma) + \xi_1)(\varphi_1(\gamma) + \xi_1 - 1)x^{\gamma, \mathbf{0}}. \quad (3.33)$$

Repeatedly applying $\rho(\bar{x}^{\varepsilon, 1_{[2]}})\rho(\bar{x}^{-\varepsilon, 1_{[2]}})$ on $\mathcal{N}$, we obtain $x^{\gamma, \mathbf{0}} \in \mathcal{N}$ by Lemma 3, (3.31), (3.32) and (3.33).

If $\varphi_1(\alpha) + \xi_1 = 0$, pick $\tau \in \text{Rad}_\phi \cap (\cap_{p \in \overline{2, 2n}} \ker_{\varphi_p}) \setminus \ker_{\varphi_1}$ such that $\varphi_1(\tau) + 1 \neq 0$.

Moreover, if there exists $q \in \overline{3, 2n}$ such that $\varphi_q(\alpha) + \xi_q \neq 0$ and $\varphi_{q'} \neq 0$, picking $\tau' \in \text{Rad}_\phi \cap (\cap_{q' \neq p \in \overline{1, 2n}} \ker_{\varphi_p}) \setminus \ker_{\varphi_{q'}}$, we have

$$\mathcal{N} \ni \bar{x}^{\tau+\tau', \mathbf{0}}.x^{\alpha, \mathbf{0}} = \epsilon_{q'} \varphi_{q'}(\tau')(\varphi_q(\alpha) + \xi_q)x^{\alpha+\tau+\tau'+\sigma_q+\varepsilon, \mathbf{0}}. \quad (3.34)$$

Since $\varphi_1(\alpha + \tau + \tau' + \sigma_q + \varepsilon) + \xi_1 \neq 0$, we go back to the previous condition from (3.31) to (3.33) with $x^{\alpha, \mathbf{0}}$ replaced by $x^{\alpha+\tau+\tau'+\sigma_q+\varepsilon, \mathbf{0}}$. So we obtain that $x^{\gamma, \mathbf{0}} \in \mathcal{N}$.

If there exists $q \in \overline{3, 2n}$ such that $\varphi_q(\alpha) + \xi_q \neq 0$ and $\varphi_{q'} \equiv 0$, then we have

$$\mathcal{N} \ni \bar{x}^{\tau, 1_{[q']}}.x^{\alpha, \mathbf{0}} = \epsilon_{q'}(\varphi_q(\alpha) + \xi_q)x^{\alpha+\tau+\sigma_q+\varepsilon, \mathbf{0}}. \quad (3.35)$$

Since $\varphi_1(\alpha + \tau + \sigma_q + \varepsilon) + \xi_1 \neq 0$, we go back to the condition from (3.31) to (3.33) with $x^{\alpha, \mathbf{0}}$ replaced by $x^{\alpha+\tau+\sigma_q+\varepsilon, \mathbf{0}}$. So we obtain that $x^{\gamma, \mathbf{0}} \in \mathcal{N}$.

If $\varphi_q(\alpha) + \xi_q = 0$ for all $q \in \overline{3, 2n}$, then there exists $\theta \in \Gamma$ such that

$$\phi(\theta, \alpha) + f(\theta) \neq 0. \quad (3.36)$$

We rechoose $\tau \in \text{Rad}_\phi \cap (\cap_{p \in \overline{2, 2n}} \ker_{\varphi_p}) \setminus \ker_{\varphi_1}$ such that $\varphi_1(\alpha + \theta + \tau + \varepsilon) + \xi_1 \neq 0$, then we have

$$\mathcal{N} \ni \bar{x}^{\theta+\tau, \mathbf{0}}.x^{\alpha, \mathbf{0}} = (\phi(\theta, \alpha) + f(\theta))x^{\alpha+\theta+\tau+\varepsilon, \mathbf{0}}. \quad (3.37)$$

We go back to the condition from (3.31) to (3.33) with $x^{\alpha, \mathbf{0}}$ replaced by $x^{\alpha+\theta+\tau+\varepsilon, \mathbf{0}}$. So we obtain that $x^{\gamma, \mathbf{0}} \in \mathcal{N}$.

Subcase 4.3. $\varphi_2(\gamma) + \xi_2 = 0$, $\varphi_1(\gamma) + 1 + \xi_1 = 0$ and there exists $q \in \overline{3, 2n}$ such that $\varphi_q(\gamma - \sigma_q) + \xi_q \neq 0$.

The proof for $x^{\gamma, \mathbf{0}} \in \mathcal{N}$ in this subcase is similar to the proof in *Subcase 4.2*. Here we omit the details.

Subcase 4.4. $\varphi_2(\gamma) + \xi_2 = 0$, $\varphi_1(\gamma) + 1 + \xi_1 = 0$ and $\varphi_q(\gamma - \sigma_q) + \xi_q = 0$ for all $q \in \overline{3, 2n}$.

Since we assumed that μ defined in Theorem 2 does not exist, there exists $\theta \in \Gamma$ such that

$$\phi(\theta, \gamma + \varepsilon - \sum_{s=1}^{n-1} \sigma_{2s+1}) + f(\theta) \neq 0. \quad (3.38)$$

Namely, we have $\phi(\gamma, \theta) - f(\theta) \neq 0$.

If $\varphi_2(\alpha) + \xi_2 \neq 0$, choose $\tau \in \text{Rad}_\phi \cap (\cap_{p \in \overline{2, 2n}} \ker_{\varphi_p}) \setminus \ker_{\varphi_1}$ such that

$$(\varphi_1(\tau + \theta) + 1)(\varphi_2(\alpha) + \xi_2) - \varphi_2(\theta)(\varphi_1(\alpha) + \xi_1) \neq 0 \quad (3.39)$$

and

$$\varphi_1(\alpha + \tau + \theta) + 1 + \xi_1 \neq 0. \quad (3.40)$$

Then we have

$$\begin{aligned}
\mathcal{N} &\ni \bar{x}^{\tau+\theta, \mathbf{0}}.x^{\alpha, \mathbf{0}} \\
&= ((\varphi_1(\tau + \theta) + 1)(\varphi_2(\alpha) + \xi_2) - \varphi_2(\theta)(\varphi_1(\alpha) + \xi_1))x^{\alpha+\tau+\theta, \mathbf{0}} \\
&\quad + (\phi(\theta, \alpha) + f(\theta))x^{\alpha+\tau+\theta+\varepsilon, \mathbf{0}} + \sum_{s=1}^{n-1} (\varphi_{2s+1}(\theta)(\varphi_{2s+2}(\alpha) + \xi_{2s+2}) \\
&\quad - \varphi_{2s+2}(\theta)(\varphi_{2s+1}(\alpha) + \xi_{2s+1}))x^{\alpha+\tau+\theta+\sigma_{2s+1}+\varepsilon, \mathbf{0}}. \tag{3.41}
\end{aligned}$$

Similarly to the discussion from (3.11) to (3.13) in *Case 1*, we obtain that $x^{\alpha+\tau+\theta, \mathbf{0}} \in \mathcal{N}$. Then we go back to *Case 3* with $x^{\alpha, \mathbf{0}}$ replaced by $x^{\alpha+\tau+\theta, \mathbf{0}}$.

Suppose $\varphi_2(\alpha) + \xi_2 = 0$ and $\varphi_1(\alpha) + \xi_1 \neq 0$. If $\varphi_2 \not\equiv 0$, choose

$$\tau \in \text{Rad}_\phi \cap (\cap_{2 \neq p \in \overline{1, 2n}} \ker \varphi_p) \setminus \ker \varphi_2 \tag{3.42}$$

such that

$$\varphi_2(\tau + \theta) \neq 0 \text{ and } \varphi_2(\alpha + \tau + \theta) + \xi_2 \neq 0. \tag{3.43}$$

Then we have

$$\begin{aligned}
\mathcal{N} &\ni \bar{x}^{\tau+\theta, \mathbf{0}}.x^{\alpha, \mathbf{0}} \\
&= (\phi(\theta, \alpha) + f(\theta))x^{\alpha+\tau+\theta+\varepsilon, \mathbf{0}} - \varphi_2(\tau + \theta)(\varphi_1(\alpha) + \xi_1)x^{\alpha+\tau+\theta, \mathbf{0}} \\
&\quad + \sum_{s=1}^{n-1} (\varphi_{2s+1}(\theta)(\varphi_{2s+2}(\alpha) + \xi_{2s+2}) - \varphi_{2s+2}(\theta)(\varphi_{2s+1}(\alpha) + \xi_{2s+1})) \\
&\quad \cdot x^{\alpha+\tau+\theta+\sigma_{2s+1}+\varepsilon, \mathbf{0}}. \tag{3.44}
\end{aligned}$$

Similarly to the discussion from (3.11) to (3.13) in *Case 1*, we obtain that $x^{\alpha+\tau+\theta, \mathbf{0}} \in \mathcal{N}$. Then we go back to *Case 3* with $x^{\alpha, \mathbf{0}}$ replaced by $x^{\alpha+\tau+\theta, \mathbf{0}}$. Assume that $\varphi_2 \equiv 0$, then we have $\mathcal{J}_2 = \mathbb{N}$ and $\xi_2 = 0$. Since we have

$$\begin{aligned}
\mathcal{N} &\ni \bar{x}^{\theta, 1[2]}.x^{\alpha, \mathbf{0}} \\
&= \sum_{s=1}^{n-1} (\varphi_{2s+1}(\theta)(\varphi_{2s+2}(\alpha) + \xi_{2s+2}) - \varphi_{2s+2}(\theta)(\varphi_{2s+1}(\alpha) + \xi_{2s+1}))x^{\alpha+\theta+\sigma_{2s+1}+\varepsilon, 1[2]} \\
&\quad + (\phi(\theta, \alpha) + f(\theta))x^{\alpha+\theta+\varepsilon, 1[2]} - (\varphi_1(\alpha) + \xi_1)x^{\alpha+\theta, \mathbf{0}}. \tag{3.45}
\end{aligned}$$

Similarly to the discussion from (3.31) to (3.33), we obtain that $x^{\alpha+\theta, \mathbf{0}} \in \mathcal{N}$. We go back to *Case 3* with $x^{\alpha, \mathbf{0}}$ replaced by $x^{\alpha+\theta, \mathbf{0}}$, then we obtain $x^{\gamma, \mathbf{0}} \in \mathcal{N}$.

Next we consider the case that $\varphi_2(\alpha) + \xi_2 = 0$, $\varphi_1(\alpha) + \xi_1 = 0$ and there exists $q \in \overline{3, 2n}$ such that $\varphi_q(\alpha) + \xi_q \neq 0$. Choose $\tau \in \text{Rad}_\phi \cap (\cap_{p \in \overline{2, 2n}} \ker \varphi_p) \setminus \ker \varphi_1$ such that $\varphi_1(\tau) + 1 \neq 0$. If $\varphi_{q'} \not\equiv 0$, picking $\tau' \in \text{Rad}_\phi \cap (\cap_{q' \neq p \in \overline{1, 2n}} \ker \varphi_p) \setminus \ker \varphi_{q'}$, we have

$$\mathcal{N} \ni \bar{x}^{\tau+\tau', \mathbf{0}}.x^{\alpha, \mathbf{0}} = \epsilon_{q'} \varphi_{q'}(\tau')(\varphi_q(\alpha) + \xi_q)x^{\alpha+\tau+\tau'+\sigma_q+\varepsilon, \mathbf{0}}. \tag{3.46}$$

Since $\varphi_2(\alpha + \tau + \tau' + \sigma_q + \varepsilon) + \xi_2 = 0$ and $\varphi_1(\alpha + \tau + \tau' + \sigma_q + \varepsilon) + \xi_1 \neq 0$, we go back to the condition from (3.44) to (3.45) with $x^{\alpha, \mathbf{0}}$ replaced by $x^{\alpha + \tau + \tau' + \sigma_q + \varepsilon, \mathbf{0}}$. If $\varphi_{q'} \equiv 0$, we have $\mathcal{J}_{q'} = \mathbb{N}$. Then we get

$$\mathcal{N} \ni \bar{x}^{\tau, 1_{[q']}}.x^{\alpha, \mathbf{0}} = \epsilon_{q'}(\varphi_q(\alpha) + \xi_q)x^{\alpha + \tau + \sigma_q + \varepsilon, \mathbf{0}}. \quad (3.47)$$

Since $\varphi_2(\alpha + \tau + \sigma_q + \varepsilon) + \xi_2 = 0$ and $\varphi_1(\alpha + \tau + \sigma_q + \varepsilon) + \xi_1 \neq 0$, we go back to the condition from (3.44) to (3.45) with $x^{\alpha, \mathbf{0}}$ replaced by $x^{\alpha + \tau + \sigma_q + \varepsilon, \mathbf{0}}$.

If $\varphi_q(\alpha) + \xi_q = 0$ for all $q \in \overline{1, 2n}$, then there exists $\theta' \in \Gamma$ such that

$$\phi(\theta', \alpha) + f(\theta') \neq 0. \quad (3.48)$$

Choose $m \in \mathbb{N}$ such that

$$\phi(m\theta' + \theta, \alpha) + f(m\theta' + \theta) \neq 0 \quad (3.49)$$

and

$$\phi(\gamma, m\theta' + \theta) - f(m\theta' + \theta) \neq 0. \quad (3.50)$$

Then we have

$$\mathcal{N} \ni \bar{x}^{m\theta' + \theta, \mathbf{0}}.x^{\alpha, \mathbf{0}} = (\phi(m\theta' + \theta, \alpha) + f(m\theta' + \theta))x^{\alpha + m\theta' + \theta + \varepsilon, \mathbf{0}}. \quad (3.51)$$

We go back to *Case 3* with $x^{\alpha, \mathbf{0}}$ replaced by $x^{\alpha + m\theta' + \theta + \varepsilon, \mathbf{0}}$, then we obtain that $x^{\gamma, \mathbf{0}} \in \mathcal{N}$.

Therefore we have proved that $x^{\gamma, \mathbf{i}} \in \mathcal{N}$ for all $\gamma \in \Gamma$ when $|\mathbf{i}| = 0$. Fixing $0 < k \in \mathbb{N}$, assume that $x^{\gamma, \mathbf{i}} \in \mathcal{N}$ for all $\gamma \in \Gamma$ with $|\mathbf{i}| < k$. Then we need to prove that $x^{\gamma, \mathbf{i}} \in \mathcal{N}$ for all $\gamma \in \Gamma$ with $|\mathbf{i}| = k$. The proof is similar to the proof for $|\mathbf{i}| = 0$. Here we omit the details.

This completes the proof of Theorem 2. $\square$

Acknowledgment

I would like to express my deep gratitude to Professor Xiaoping Xu for all his advice, instructions and encouragements.

References

- [1] D. Dokovic and K. Zhao, Generalized Cartan type W Lie algebras in characteristic 0, *J. Algebra* **195** (1997), 170-210.
- [2] V. G. Kac, A description of filtered Lie algebras whose associated graded Lie algebras are of Cartan types, *Math. USSR-Izvestija* **8** (1974), 801-835.

- [3] N. Kawamoto, Generalizations of Witt algebras over a field of characteristic zero, *Hiroshima Math. J.* **16** (1986), 417-462.
- [4] T. Larsson, Conformal fields: A class of representations of $\text{Vect}(\mathbb{N})[J]$, *Internat. J. Modern Phys. A* **7** (1992), no. 26, 6493-6508.
- [5] W. Lin and S. Tan, Representations of the Lie algebra for quantum torus, *J. Algebra* **275** (2004), 250-274.
- [6] J. M. Osborn, New simple infinite-dimensional Lie algebras of characteristic 0, *J. Algebra* **185** (1996), 820-835.
- [7] J. M. Osborn and K. Zhao, Generalized Poisson brackets and Lie algebras for type H in characteristic 0, *Math. Z.* **230** (1999), 107-143.
- [8] D. P. Passman, Simple Lie algebras of Witt type, *J. Algebra* **206** (1998), 682-692.
- [9] S. E. Rao, Irreducible representations of the Lie algebra of the diffeomorphisms of a d-dimensional torus, *J. Algebra* **182** (1996), 401-421.
- [10] S. E. Rao, Representations of Witt algebras, *Publ. Rims, Kyoto Univ.* **29** (1994), 191-201.
- [11] G. Shen, Graded modules of graded Lie algebras of Cartan type (I)–mixed product of modules, *Sci. China Ser. A* **29** (1986), 570-581.
- [12] G. Shen, Graded modules of graded Lie algebras of Cartan type (II)–positive and negative graded modules, *Sci. China Ser. A* **29** (1986), 1009-1019.
- [13] G. Shen, Graded modules of graded Lie algebras of Cartan type (III)–irreducible modules, *Chinese Ann. Math. Ser. B* **9** (1988), 404-417.
- [14] Y. Su, Poisson brackets and structure of nongraded Hamiltonian Lie algebras related to locally-finite derivations, *Canad. J. Math.* **55** (2003), 856-896.
- [15] Y. Su and X. Xu, Structure of divergence-free Lie algebras, *J. Algebra* **243** (2001), 557-595.
- [16] Y. Su and X. Xu, Structure of contact Lie algebras related to locally-finite derivations, *Manuscripta Mathematica* **112** (2003), 231-255.
- [17] Y. Su and X. Xu, Central simple Poisson algebras, *Science in China Series A* **47** (2004), no. 2, 245-263.
- [18] Y. Su, X. Xu and H. Zhang, Derivation-simple algebras and the structures of Lie algebras of Witt type, *J. Algebra* **233** (2000), 642-662.
- [19] Y. Su and J. Zhou, Some representations of nongraded Lie algebras of generalized Witt type, *J. Algebra* **246** (2001), 721-738.

- [20] Y. Su and K. Zhao, Generalized Virasoro and super-Virasoro algebras and modules of intermediate series, *J. Algebra* **252** (2002), 1-19.
- [21] X. Xu, Generalizations of the Block algebras, *Manuscripta Math.* **100** (1999), 489-518.
- [22] X. Xu, New generalized simple Lie algebras of Cartan type over a field with characteristic 0, *J. Algebra* **224** (2000), 23-58.
- [23] X. Xu, Quadratic conformal superalgebras, *J. Algebra* **231** (2000), 1-38.
- [24] X. Xu, Equivalence of conformal superalgebras to Hamiltonian superoperators, *Algebra Collquium* **8** (2001), 63-92.
- [25] K. Zhao, Generalized Cartan type S Lie algebras in characteristic zero, II, *Pacific J. Math.* **192** (2000), 431-454.
- [26] K. Zhao, Weight modules over generalized Witt algebras with 1-dimensional weight spaces, *Forum Math.* **16** (2004), 725-748.
- [27] Y. Zhao, Irreducible representations of nongraded Witt type Lie algebras, *J. Algebra* **298** (2006), 540-562.